

RESEARCH

Open Access

# Structure-activity-relationship study of *N*-acyl-*N*-phenylpiperazines as potential inhibitors of the Excitatory Amino Acid Transporters (EAATs): improving the potency of a micromolar screening Hit is not truism

Tri HV Huynh, Charles S Demmer, Bjarke Abrahamsen, Emil Marcher, Mikael Frykman, Anders A Jensen and Lennart Bunch\*

## Abstract

The excitatory amino acid transporters (EAATs) are transmembrane proteins responsible for the uptake of (*S*)-glutamate from the synaptic cleft. To date, five subtypes EAAT1–5 have been identified for which selective inhibitors have been discovered for EAAT1 and EAAT2. By screening of a commercially available compound library consisting of 4,000 compounds, *N*-acyl-*N*-phenylpiperazine analog ( $\pm$ )-*exo*-1 was identified to be a non-selective inhibitor at EAAT1–3 displaying  $IC_{50}$  values in the mid-micromolar range (10  $\mu$ M, 40  $\mu$ M and 30  $\mu$ M at EAAT1, 2 and 3, respectively). Subsequently, we designed and synthesized a series of analogs to explore the structure-activity-relationship of this scaffold in the search for analogs characterized by increased inhibitory potency and/or EAAT subtype selectivity. Despite extensive efforts, all analogs of ( $\pm$ )-*exo*-1 proved to be either inactive or to have least 3-fold lower inhibitory potency than the lead, and furthermore none of the active analogs displayed selectivity for a particular subtype amongst the EAAT1–3. On the basis of our findings, we speculate that ( $\pm$ )-*exo*-1 binds to a recess (deepening) on the EAAT proteins than a well-defined pocket.

**Keywords:** Excitatory amino acid transporters, EAATs, Rational ligand design, Medicinal chemistry

## Background

In the central nervous system (CNS), the excitatory amino acid transporters (EAATs) are transmembrane proteins responsible for the uptake of (*S*)-glutamate (Glu) from the synaptic cleft. Five subtypes have been identified, named EAAT1–EAAT5 in humans and GLAST, GLT-1, EAAC1, EAAT4 and EAAT5, respectively, in rodents. (Bunch et al. 2009) While EAAT5 is found exclusively in the retina, subtypes EAAT1–4 are expressed differentially within the CNS with respect to brain regions as well as at the cellular level: EAAT1 and EAAT2 are expressed primarily on astrocytes, but EAAT2 is also found in neurons, astrocytes and oligodendrocytes. (Lauriat et al. 2007) Subtype EAAT3 is


distributed predominantly in postsynaptic neuronal sites, (Nieoullon et al. 2006) whereas EAAT4 is distributed in Purkinje cells as well as in the cerebral cortex. (Massie et al. 2001) Discovery of subtype selective ligands for the EAATs has attracted much attention over the past decade, (Jensen et al. 2009) the latest being the disclosure of UCPH-101 as first subtype selective EAAT1-inhibitor (Figure 1). (Jensen et al. 2009; Erichsen et al. 2010; Huynh et al. 2012,ab).

## Results and discussion

From screening of a 4,000 compounds-library at HEK293 cells stably expressing human EAAT1–3 *N*-acyl-*N*-phenylpiperazine analog ( $\pm$ )-*exo*-1 was identified as a non-selective inhibitor at the transporter exhibiting  $IC_{50}$  values in the mid-micromolar range (10  $\mu$ M, 40  $\mu$ M and 30  $\mu$ M at EAAT1, -2 and -3, respectively, Figure 2).

\* Correspondence: lebu@sund.ku.dk

Department of Drug Design and Pharmacology, Faculty of Health and Medical Sciences, University of Copenhagen, Universitetsparken 2, Copenhagen, OE 2100, Denmark




**Figure 1** Chemical structures of the EAAT1-selective inhibitor UCPH-101, (Jensen et al. 2009; Erichsen et al. 2010) and EAAT2-selective inhibitors DHK, (Jensen & Bräuner-Osborne 2004) DPAG, (Sagot et al. 2008) and WAY-213613 (Dunlop et al. 2005).

Although phenylpiperazines are promiscuous hits in high-throughput screenings (HTS) and a frequent core skeleton in marketed drugs, (Millan et al. 2001; Fragasso et al. 2006; Weisberg et al. 2007) we were motivated to explore the structure-activity-relationship (SAR) of this new class of EAAT inhibitors. A conventional medicinal chemistry analysis of ( $\pm$ )-*exo*-1 suggests that the amide functionality, the aniline nitrogen, the phenyl ring and the trifluoromethyl group may play key roles in binding of this class of EAAT inhibitors. Consequently, the chemical structure can be broken down into three fragments: a core skeleton being the acyl-phenylpiperazine scaffold and two substituents being the trifluoromethyl- and the bicyclo [2.2.1]heptanyl group (Figure 1). The SAR study was designed as to study the influence of one of the two substituents (Figure 1) individually including the stereochemical organization around the  $\alpha$ -carbonyl carbon.













The SAR study commenced by investigation of the influence of bicycle[2.2.1]heptanyl group on the EAAT inhibitory activity. The stereochemical configuration of the  $\alpha$ -carbonyl carbon was addressed by the synthesis of *endo*-conformer ( $\pm$ )-*endo*-1 (Table 1) from commercially available ( $\pm$ )-*endo*-carboxylic acid and *N*-(3-trifluoromethylphenyl)-piperazine **4**, the latter prepared by a palladium-catalyzed amination of commercially available piperazine (Scheme 1). (Nishiyama et al. 1998) Furthermore, the racemic and diastereomeric mixture ( $\pm$ )-*endo*-*exo*-1 was prepared from the corresponding acid ( $\pm$ )-*endo*-*exo*-6 obtained from

oxidation of commercially available ( $\pm$ )-*endo*-*exo*-bicyclo [2.2.1]heptanymethyl alcohol (( $\pm$ )-*endo*-*exo*-5) (Scheme 2) using  $\text{KMnO}_4$  and  $\text{K}_2\text{CO}_3$  in  $\text{H}_2\text{O}$ . Isolation of ( $\pm$ )-*endo*-*exo*-6 turned out to be difficult for which reason it was used directly in the next step. (Gudipati et al. 1993) To search for the optimal bulkiness of the lipophilic substituent, larger as well as smaller rigid hydrophobic ring-systems were introduced (**2.1–2.5**). Moreover, analogs **2.6–2.10** comprising alkyl group of varying length and bulkiness were designed to explore the effect of increased flexibility of this substituent on ligand binding. Furthermore, analogs **2.11–2.17** address if the bicyclo [2.2.1]heptane could be substituted for an aromatic moiety, whereas the analogs **2.18–2.21** were designed to explore the distinct substitution for a hydrophobic group. The synthesis of piperazine analogs **1** and **2.1–2.21** was carried out by amidation of **4** using the respective carboxylic acids, acid chlorides, benzenesulfonyl chloride and benzyl carbonochloridate afforded the corresponding amides in moderate to good yields (Scheme 1). The rationally designed 3-trifluoromethylphenylpiperazine analogs were supplemented by commercially available analogs **2.22–2.31**, as a quick way of expanding the SAR into the chemical space beyond rational guidance. Finally, the importance of the amide functionality was explored by the synthesis of carbamate **2.32** by acylation of **4** with carbonochloridate, sulfonamide **2.33** by treatment of **4** with phenylsulfonyl chloride, amine **2.34** by reduction of ( $\pm$ )-*endo*-*exo*-1 with
















**Figure 2** Chemical structure of screening hit *N*-acyl-*N*-phenylpiperazine ( $\pm$ )-*exo*-1 displaying inhibitory activity at EAAT1–3 in the mid-micromolar range ( $\text{IC}_{50}$  = 10, 40 and 30  $\mu\text{M}$ , respectively).













**Table 1 Pharmacological characterization of piperazine analogs 1 and 2.1–2.40 as inhibitors at HEK293 cells stably expressing human EAAT1-3 in the [<sup>3</sup>H]-D-aspartate uptake assay (Jensen & Bräuner-Osborne 2004)**

				
	R <sup>1</sup>	EAAT1	EAAT2	EAAT3
(±)- <i>exo</i> -1		10 [5.03 ± 0.12]	40 [4.44 ± 0.14]	30 [4.78 ± 0.10]
(±)- <i>endo</i> -1		14 [4.88 ± 0.09]	32 [4.52 ± 0.12]	10 [5.04 ± 0.13]
(±)- <i>endo-exo</i> -1		14 [4.87 ± 0.08]	42 [4.42 ± 0.11]	14 [4.94 ± 0.16]
2.1		>300	>300	>300
2.2		>300	>300	>300
2.3		~100	~100	~100
2.4		>300	>300	>300
2.5		>300	>300	>300
2.6		>1000	>300	>300
2.7		>1000	>100	>100
2.8		>300	>300	>300

**Table 1 Pharmacological characterization of piperazine analogs 1 and 2.1–2.40 as inhibitors at HEK293 cells stably expressing human EAAT1-3 in the [<sup>3</sup>H]-D-aspartate uptake assay (Jensen & Bräuner-Osborne 2004) (Continued)**

2.9		>300	>300	>300
2.10		>300	>300	>300
2.11		~150	>100	>100
2.12		>100	>100	>100
2.13		>300	>300	>300
2.14		>300	>300	>300
2.15		>300	>300	>300
2.16		>100	>100	>100
2.17		>300	>300	>1000
2.18		>300	>300	>300
2.19		>300	>300	>300
2.20		>300	>300	>300
2.21		>300	>300	>300


**Table 1 Pharmacological characterization of piperazine analogs 1 and 2.1–2.40 as inhibitors at HEK293 cells stably expressing human EAAT1-3 in the [ $^3$ H]-D-aspartate uptake assay (Jensen & Bräuner-Osborne 2004) (Continued)**


2.22		~100	>100	>100
2.23		>300	>300	>100
2.24		>100	>100	>100
2.25		>300	>100	>300
2.26		>300	>300	>300
2.27		>300	>300	>300
2.28		>300	>300	>300
2.29		>300	>100	>100
2.30		>300	>1000	>300
2.31		>100	>100	>100
2.32		>1000	>300	>300
2.33		>1000	>300	>1000

**Table 1 Pharmacological characterization of piperazine analogs 1 and 2.1–2.40 as inhibitors at HEK293 cells stably expressing human EAAT1-3 in the [<sup>3</sup>H]-D-aspartate uptake assay (Jensen & Bräuner-Osborne 2004) (Continued)**

(±)-endo-exo-2.34		>300	>300	>300
2.35		>300	>300	>1000
2.36		>300	>100	>300
2.37		>300	>100	>100
2.38		>300	>300	>300
2.39		>100	>100	>100
2.40		>500	>1000	>500

All values are given as IC<sub>50</sub> in μM with pIC<sub>50</sub> ± S.E.M. values in brackets (for the active analogs).





$\text{LiAlH}_4$ , (Cook et al. 1992) and *N*-benzyl analog **2.35** by alkylation of phenylpiperazine **4**. (Burkhard et al. 2010) In addition, these analogs were supplemented by five commercially available structurally diverse analogs **2.36–2.40**.

We then turned to the design of analogs for investigation of the influence on EAAT inhibitory activity of the chemical nature of the trifluoromethyl group as well as its position on the phenyl ring. A series of 12 analogs were included in the SAR study, all wherein the (±)-endo-exo-bicyclic [2.2.1]-acyl group was conserved (analog **3.1–3.12**, Table 2): Simplification of the chemical structure by depletion of the trifluoromethyl group provides analog (±)-endo-exo-3.1, while shifting the 3-trifluoromethyl group to the 4- or 2-positions affords analogs (±)-endo-exo-3.2 and (±)-endo-exo-3.5, respectively (Table 2 and Scheme 2). The latter two analogs were supplemented by commercially available analogs (±)-endo-exo-3.3, (±)-endo-exo-3.4 and (±)-endo-exo-3.6. Continuing the design stage, substitution of the 3-trifluoromethyl group for a chloride, hydroxyl-, cyano- and methoxy group, provides analogs (±)-endo-exo-3.7–3.10 respectively (Table 2), whereas 2,4-difluorophenyl analog (±)-endo-exo-3.11 was included due to readily available starting materials. Analog (±)-endo-exo-3.12 could be obtained from commercial suppliers and thus included with the notion that it comprises an *N*-diphenylmethyl group, which is indeed chemically distinct from the *N*-3-trifluoromethylphenyl group and furthermore the connecting nitrogen will be protonated at physiological pH=7.4. The analogs were synthesized starting from the appropriate phenylpiperazine and (±)-endo-exo-6 under standard coupling conditions (TBTU, DIPEA in DMF) for the amide formation to afford the target compounds in moderate yields (Scheme 3) (Balalaie et al. 2007).

### Pharmacological characterization

In total, 54 piperazine analogs **2.1–2.40** and **3.1–3.12** were characterized pharmacologically at stable HEK293 cells expressing human EAAT1–3 in a [ $^3\text{H}$ ]-D-aspartate uptake assay, (Jensen & Bräuner-Osborne 2004) and the results are summarized in Table 1 and Table 2. The *endo*-isomer **1** (*endo*-isomer) displayed inhibitory activities at EAAT1–3 comparable with those of the *exo*-isomer **1** (lead structure) ( $\text{IC}_{50}$  = 14  $\mu\text{M}$ , 32  $\mu\text{M}$  and 10  $\mu\text{M}$  vs.

10  $\mu\text{M}$ , 40  $\mu\text{M}$  and 30  $\mu\text{M}$ , respectively). In line with this, *endo-exo* **1**, which is a 1:1 ratio of *endo/exo* moiety displayed  $\text{IC}_{50}$  values at EAAT1–3 of  $\text{IC}_{50}$  = 14  $\mu\text{M}$ , 42  $\mu\text{M}$  and 14  $\mu\text{M}$ , respectively. Usually, such findings would lead to the conclusion that the bicyclo-[2.2.1]-heptanyl group occupies a promiscuous hydrophobic pocket, which could be optimized for increased potency. However, upon increasing or decreasing the hydrophobic bulk and/or flexibility, a clear drop in potency was observed (analog **2.1–2.17**, Table 1). Except for analogs **2.3** and **2.11**, which displayed only a 5–15 fold drop in inhibitory potency across the subtypes, all of these analogs would be characterized as *inactive* ( $\text{IC}_{50}$  >100  $\mu\text{M}$  or >300  $\mu\text{M}$ , Table 1). These findings could open up for the hypothesis that the pocket is indeed not hydrophobic but instead hydrophilic in nature. Upon binding of the hydrophobic alkane group, water molecules are forced out and ligand binding is entropically driven rather than enthalpically. Thus analogs **2.18–2.31**, which comprise a hydrophobic group, could be potential inhibitors. However, none of these displayed any inhibitory activity at EAAT1–3 ( $\text{IC}_{50}$  >100, >300 or >1000  $\mu\text{M}$ , Table 1). Continuing the characterization, neither the carbamate **2.32** nor the sulfonamide **2.33** analog displayed inhibitory activity at EAAT1–3, and likewise all amines **2.34–2.40** were found to be inactive at all three subtypes.

The pharmacological results for the twelve analogs **3.1–3.12**, which address the influence of substituent 2 (Figure 2) on inhibitory activity at EAAT1–3 are summarized in Table 2. While it was not surprising that removal or repositioning of the 3-trifluoromethyl group (analog **3.1**, **3.2** and **3.5**, respectively) resulted in loss of inhibitory activity, the further nine analogs **3.3**, **3.4**, **3.6–3.12** were also inactive or displayed at least a 3-fold lower inhibitory activity than (±)-*exo*-**1**.

### Conclusion


In conclusion, screening of a compound library identified (±)-*exo*-**1** as a broad range EAAT1–3 inhibitor exhibiting  $\text{IC}_{50}$  values at the three transporters in the mid-micromolar range. Subsequently, rational design and synthesis of 33 analogs of (±)-*exo*-**1** was carried out, together with the purchase of 21 analogs. Thus, a total of 54 piperazine analogs were characterized pharmacologically as inhibitors

**Table 2 Pharmacological characterization of analogs 3.1–3.12 as inhibitors at HEK293 cells stably expressing human EAAT1–3 in the [<sup>1</sup>H]-D-aspartate uptake assay (Jensen & Bräuner-Osborne 2004)**

	R <sup>2</sup>	EAAT1	EAAT2	EAAT3
(±)-endo-exo-3.1		>300	>300	>300
(±)-endo-exo-3.2		>300	>300	>300
(±)-endo-exo-3.3		>100	>100	>100
(±)-endo-exo-3.4		>100	~100	>300
(±)-endo-exo-3.5		~300	>300	>300
(±)-endo-exo-3.6		>100	~100	>100
(±)-endo-exo-3.7		~300	~300	>300
(±)-endo-exo-3.8		>300	>300	>300
(±)-endo-exo-3.9		>300	>300	>300
(±)-endo-exo-3.10		>300	>300	>300
(±)-endo-exo-3.11		>300	>300	>300
(±)-endo-exo-3.12		>100	~100	>100

All values are given as IC<sub>50</sub> in μM with pIC<sub>50</sub> ± S.E.M. values in brackets (for the active analogs).





**Scheme 3** Synthesis of piperazine analogs 3.1, 3.5, 3.8–3.11 by use of the coupling reagent TBTU <sup>a</sup>. <sup>a</sup> Reagents and conditions: (a) TBTU, DIPEA, dry DMF, rt, 20 h, 40–58%.

at EAAT1–3 but only the *endo* diastereomer (±)-*endo*-1 displayed inhibitory potency in the mid-micromolar range comparable to that of the lead structure (±)-*exo*-1. The remaining analogs were inactive or at least three fold weaker inhibitors at EAAT1–3 than the lead, none of them displaying signs of subtype-selectivity. Given the structural diversity of the analogs characterized pharmacologically, we speculate if the lead structure (±)-*exo*-1 adheres to a recess (deepening) in the surface of the protein rather than binds in an organized way to a well-defined pocket.

#### Experimental section

All commercially available reagents were used without further purification. THF was distilled over sodium/benzophenone, Et<sub>2</sub>O was dried over neatly cut sodium and dichloromethane was dried over 3 Å molecular sieves. All solvents were tested for water content using a Karl Fisher apparatus. All reactions involving dry solvents or sensitive agents were performed under a nitrogen atmosphere, and glassware was dried prior to use. All reactions were monitored by analytical thin-layer chromatography (TLC, Merck silica gel 60 F<sub>254</sub> aluminum sheets). Flash chromatography was carried out using Merck silica gel 60A (35–70 micron). <sup>1</sup>H NMR spectra were recorded on a 300 MHz Varian Mercury 300BB or a 400 MHz Avance Bruker and <sup>13</sup>C NMR spectra on a 75 MHz Varian Gemini 2000BB or a 100 MHz Avance Bruker. Preparative HPLC was done using Agilent Prep HPLC systems with Agilent 1100 series pump, Agilent 1200 series diode array, multiple wavelength detector (G1365B), and Agilent PrepHT High Performance Preparative Cartridge Column (Zorbax, 300 SB-C18 Prep HT, 21.2 × 250 mm, 7 μm). LC-MS spectra were recorded using an Agilent 1200 series solvent delivery system equipped with an autoinjector coupled to an Agilent 6400 series triple quadrupole mass spectrometer equipped with an electrospray ionization source. Gradients of 5% aqueous MeCN + 0.05% HCOOH (eluent A), and 95% aqueous MeCN + 0.043% formic acid (eluent B) were employed. Melting points were measured using a MPA 100 Optimelt automatic melting point system and are stated

uncorrected. Compounds were dry either under high vacuum or freeze dried using a Holm & Halby, Heto LyoPro 6000 freeze-drier.

#### General procedure a: synthesis of amides using *O*-benzotriazole-1-yl-*N,N,N',N'*-tetramethyluronium tetrafluoroborate (TBTU) as coupling reagent

To a suspension of the appropriate phenylpiperazine analog (0.33 mmol), the carboxylic acid (0.40 mmol) and TBTU (0.43 mmol) in dry DMF (4 mL) under an N<sub>2</sub> atmosphere, was added DIPEA (1.32 mmol) and reaction mixture was stirred for 20 h at rt. The reaction mixture was quenched with brine (5 mL) and extracted with dichloromethane (3 × 20 mL). The combined organic phases were washed with H<sub>2</sub>O (20 mL) and brine (20 mL) and dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. After concentration *in vacuo*, the crude product was purified by column chromatography on silica gel in accordance with details described for the analog.

#### General procedure B: synthesis of amides using acid chlorides

To a suspension of 1-(3-(trifluoromethyl)phenyl)piperazine (4) (0.33 mmol) in dry dichloromethane (5 mL) at 0°C under a N<sub>2</sub> atmosphere was added Et<sub>3</sub>N (0.91 mmol). The reaction mixture was stirred for 10 min at 0°C, then the appropriate acid chloride (0.48 mmol) was added and stirring continued for 30 minutes at rt. The reaction mixture was quenched with sat. NH<sub>4</sub>Cl (5 mL) and extracted with dichloromethane (3 × 20 mL). The combined organic phases were washed with H<sub>2</sub>O (20 mL) and brine (20 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. After concentration *in vacuo*, the crude product was purified by column chromatography on silica gel in accordance with details described for the analog.

#### (±)-endo-Bicyclo[2.2.1]heptan-2-yl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone ((±)-endo-1)

Obtained from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and commercially available (±)-endo-bicyclo[2.2.1]heptane-2-carboxylic acid by general procedure A in

61% yield as a pale-yellow oil.  $R_f$  0.25 (heptane/EtOAc 3:1).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.36 (t,  $J$  = 8.4 Hz, 1H), 7.12–7.04 (m, 3H), 3.94–3.89 (m, 1H), 3.79–3.62 (m, 3H), 3.30–3.08 (m, 4H), 2.95 (dt,  $J$  = 10.8, 4.2 Hz, 1H), 2.40 (br s, 1H), 2.29 (br s, 1H), 1.95 (ddd,  $J$  = 12.0, 4.5, 2.4 Hz, 1H), 1.64–1.28 (m, 7H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  172.3, 151.1, 131.5 (q,  $J$  = 31.5 Hz), 129.7, 124.2 (q,  $J$  = 270.7 Hz), 119.3, 116.5 (q,  $J$  = 3.8 Hz), 112.6 (q,  $J$  = 3.8 Hz), 49.5, 45.2, 43.7, 41.7, 40.8, 40.5, 40.2, 37.1, 32.2, 29.1, 21.1. LC-MS ( $m/z$ ) calcd for  $\text{C}_{19}\text{H}_{23}\text{F}_3\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 353.2; found, 353.2. Anal. calcd for  $\text{C}_{19}\text{H}_{23}\text{F}_3\text{N}_2\text{O} \times \text{HCl}$ : C 58.69, H 6.22, N 7.20 found C 59.28, H 6.24, N 7.18.

**( $\pm$ )-endo-exo-Bicyclo[2.2.1]heptan-2-yl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (( $\pm$ )-endo-exo-1)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and ( $\pm$ )-endo-exo-bicyclo[2.2.1]heptane-2-carboxylic acid (( $\pm$ )-endo-exo-(6)) by general procedure A in 36% yield as a pale-yellow oil.  $R_f$  0.50 (heptane/EtOAc 1:1).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.38–7.33 (m, 1H), 7.12–7.02 (m, 3H), 3.98–3.96 (m, 1H), 3.75 (br s, 2H), 3.67 (br s, 2H), 3.21 (br s, 4H), 2.98–2.92 (m, 0.6H), 2.40 (br s, 1.4H), 2.29 (br s, 1H), 1.95–1.92 (m, 1H), 1.62–1.25 (m, 6H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  173.1, 151.3, 131.5 (q,  $J$  = 31.5 Hz), 129.7, 124.2 (q,  $J$  = 270.7 Hz), 119.2, 116.5 (q,  $J$  = 3.8 Hz), 112.7 (q,  $J$  = 3.8 Hz), 50.2, 49.8, 45.2, 43.7, 41.7, 40.8, 40.4, 36.7, 32.3, 28.9, 21.9. LC-MS ( $m/z$ ) calcd for  $\text{C}_{19}\text{H}_{23}\text{F}_3\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 353.2; found, 353.4. HPLC: purity<sub>254</sub> > 99%.

**Bicyclo[2.2.2]octan-2-yl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.1)**

Obtained from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and racemic ( $\pm$ )-endo-exo-bicyclo[2.2.2]octane-2-carboxylic acid (( $\pm$ )-endo-exo-(6)) by general procedure A in 59% yield as a pale-yellow oil.  $R_f$  0.95 (1:1 heptane/EtOAc).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.12 (t,  $J$  = 6.0 Hz, 1H), 6.50 (dd,  $J$  = 6.0, 1.0 Hz, 1H), 6.42–6.35 (m, 2H), 3.82 (br s, 2H), 3.64 (br s, 2H), 3.32 (br s, 4H), 2.79–2.76 (m, 1H), 2.19–2.16 (m, 1H), 1.79–1.73 (m, 1H), 1.69–1.18 (m, 10H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  174.0, 151.2, 131.5 (q,  $J$  = 31.5 Hz), 129.7, 125.6 (q,  $J$  = 270.7 Hz), 119.3, 116.5 (q,  $J$  = 3.8 Hz), 112.6 (q,  $J$  = 3.8 Hz), 49.2, 45.2, 41.6, 38.8, 28.4, 27.6, 26.5, 25.3, 25.2, 23.9, 21.5. LC-MS ( $m/z$ ) calcd for  $\text{C}_{20}\text{H}_{25}\text{F}_3\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 366.2; found, 366.2. HPLC: purity<sub>254</sub> > 98%.

**Adamantan-1-yl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.2)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and 1-adamantanecarboxylic acid by general procedure A in 73% yield as a pale-yellow oil.  $R_f$  0.68 (100% EtOAc).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.39–7.32 (m, 1H), 7.12–7.04

(m, 3H), 3.87 (dd,  $J$  = 6.0, 3.0 Hz, 4H), 3.21 (dd,  $J$  = 6.0, 3.0 Hz, 4H), 2.06–2.03 (m, 9H), 1.70–1.77 (m, 6H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  176.2, 151.5, 131.1 (q,  $J$  = 31.5 Hz), 130.1, 127.2 (q,  $J$  = 270.8 Hz), 119.4, 116.9 (q,  $J$  = 3.8 Hz), 112.8 (q,  $J$  = 3.8 Hz), 55.2, 49.7, 49.6, 45.4, 43.9, 42.1, 39.5, 37.1, 28.9, 19.1. LC-MS ( $m/z$ ) calcd for  $\text{C}_{22}\text{H}_{27}\text{F}_3\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 393.2; found, 393.2. Anal. calcd for  $\text{C}_{22}\text{H}_{27}\text{F}_3\text{N}_2\text{O} \times \text{HCl}$ : C 61.61, H 6.58, N 6.53 found C 61.38, H 6.38, N 6.38.

**Cyclohexyl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.3)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and cyclohexanecarboxylic acid by general procedure A in 59% yield as a yellow oil.  $R_f$  0.15 (heptane/EtOAc 3:1).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.35 (t,  $J$  = 7.8 Hz, 1H), 7.12–7.03 (m, 3H), 3.78 (br s, 2H), 3.67 (br s, 2H), 3.21 (br s, 4H), 2.50 (tt,  $J$  = 11.4, 3.3 Hz, 1H), 1.83–1.69 (m, 5H), 1.55 (dq,  $J$  = 11.4, 3.9 Hz, 2H), 1.35–1.23 (m, 3H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  175.0, 151.5, 131.9 (q,  $J$  = 31.5 Hz), 130.1, 124.6 (q,  $J$  = 270.8 Hz), 119.7 (q,  $J$  = 1.5 Hz), 117.0 (q,  $J$  = 3.8 Hz), 113.0 (q,  $J$  = 3.8 Hz), 49.8, 49.5, 45.1, 40.9, 29.8, 26.3. LC-MS ( $m/z$ ) calcd for  $\text{C}_{18}\text{H}_{23}\text{F}_3\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 341.1; found, 341.1. Anal. calcd for  $\text{C}_{18}\text{H}_{23}\text{F}_3\text{N}_2\text{O} \times \text{HCl}$ : C 57.37, H 6.42, N 7.43 found C 57.38, H 6.20, N 7.38.

**Cyclopentyl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.4)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and cyclopentanecarbonyl chloride by general procedure B in 85% yield as a yellow oil.  $R_f$  0.50 (heptane/EtOAc 1:1).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.37 (t,  $J$  = 8.0 Hz, 1H), 7.13–7.06 (m, 3H), 3.80 (dd,  $J$  = 8.0, 4.0 Hz, 2H), 3.70 (dd,  $J$  = 8.0, 4.0 Hz, 2H), 3.24 (dd,  $J$  = 8.0, 4.0 Hz, 2H), 3.19 (dd,  $J$  = 8.0, 4.0 Hz, 2H), 1.92–1.54 (m, 9H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  176.1, 147.0, 134.2 (q,  $J$  = 31.5 Hz), 132.4, 125.0 (q,  $J$  = 270.7 Hz), 124.9, 118.1 (q,  $J$  = 3.8 Hz), 117.6 (q,  $J$  = 3.8 Hz), 54.7, 43.1, 42.4, 31.6, 27.5. LC-MS ( $m/z$ ) calcd for  $\text{C}_{17}\text{H}_{21}\text{F}_3\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 327.2; found, 327.2. HPLC: purity<sub>254</sub> > 97%.

**Cyclopropyl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.5)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and cyclopropanecarboxylic acid by general procedure A in 51% yield as a yellow oil.  $R_f$  0.55 (100% EtOAc).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.39–7.32 (m, 1H), 7.12–7.05 (m, 3H), 3.83 (br s, 4H), 3.26 (br s, 2H), 3.24 (br s, 2H), 1.78 (m, 1H), 1.05–1.00 (m, 2H), 0.84–0.78 (m, 2H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  172.5, 151.5, 131.9 (q,  $J$  = 31.5 Hz), 130.1, 127.5 (q,  $J$  = 270.8 Hz), 119.6 (q,  $J$  = 0.8 Hz), 116.9 (q,  $J$  = 3.8 Hz), 113.0 (q,  $J$  = 3.8 Hz), 49.5, 49.3, 45.6, 42.2, 11.4, 8.0. LC-MS ( $m/z$ ) calcd for  $\text{C}_{15}\text{H}_{17}\text{F}_3\text{N}_2\text{O}$

[M+H<sup>+</sup>], 299.1; found, 299.1. Anal. calcd for C<sub>15</sub>H<sub>17</sub>F<sub>3</sub>N<sub>2</sub>O × 1HCl: C 53.82, H 5.42, N 8.37 found C 56.15, H 5.15, N 8.17.

**2,2-Dimethyl-1-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)propan-1-one (2.6)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and pivaloyl chloride by general procedure B in 78% yield as a clear oil. *R*<sub>f</sub> 0.28 (heptane/EtOAc 3:1). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.35 (dt, *J* = 8.1, 0.9 Hz, 1H), 7.11–7.03 (m, 3H), 3.81 (dd, *J* = 5.1, 5.1 Hz, 4H), 3.21 (dd, *J* = 5.1, 5.1 Hz, 4H), 1.32 (s, 9H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 176.4, 151.1, 131.5 (q, *J* = 31.5 Hz), 129.7, 124.2 (q, *J* = 270.8 Hz), 119.1 (q, *J* = 1.5 Hz), 116.5 (q, *J* = 3.8 Hz), 112.4 (q, *J* = 3.8 Hz), 49.0, 44.8, 38.7, 28.4. LC-MS (*m/z*) calcd for C<sub>16</sub>H<sub>21</sub>F<sub>3</sub>N<sub>2</sub>O [M+H<sup>+</sup>], 315.2; found, 315.2. Anal. calcd for C<sub>16</sub>H<sub>21</sub>F<sub>3</sub>N<sub>2</sub>O × 1HCl: C 55.11, H 6.21, N 7.96 found C 54.78, H 6.32, N 7.99.

**1-(4-(3-(Trifluoromethyl)phenyl)piperazin-1-yl)ethanone (2.7)**

Obtained from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and acetic acid by general procedure A in 64% yield as a yellow oil. *R*<sub>f</sub> 0.22 (100% EtOAc). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.36 (t, *J* = 9.0 Hz, 1H), 7.13–7.04 (m, 3H), 3.78 (dd, *J* = 6.0, 3.0 Hz, 2H), 3.64 (dd, *J* = 6.0, 3.0 Hz, 2H), 3.24 (dd, *J* = 6.0, 3.0 Hz, 2H), 3.20 (dd, *J* = 6.0, 3.0 Hz, 2H), 2.15 (s, 3H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 169.4, 151.4, 131.9 (q, *J* = 31.5 Hz), 130.1, 124.6 (q, *J* = 270.8 Hz), 119.7 (q, *J* = 1.5 Hz), 117.1 (q, *J* = 3.8 Hz), 113.1 (q, *J* = 3.8 Hz), 49.5, 49.3, 46.4, 41.6, 21.8. LC-MS (*m/z*) calcd for C<sub>13</sub>H<sub>15</sub>F<sub>3</sub>N<sub>2</sub>O [M+H<sup>+</sup>], 273.1; found, 273.1. Anal. calcd for C<sub>13</sub>H<sub>15</sub>F<sub>3</sub>N<sub>2</sub>O × 1HCl: C 50.58, H 5.22, N 9.07 found C 50.73, H 5.02, N 8.74.

**3-Methyl-1-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)butan-1-one (2.8)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and isovaleryl chloride by general procedure B in 59% yield as a pale-yellow oil. *R*<sub>f</sub> 0.40 (heptane/EtOAc 1:1). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.39 (t, *J* = 6.9 Hz, 1H), 7.16–7.10 (m, 3H), 3.82 (dd, *J* = 6.0, 3.0 Hz, 2H), 3.68 (dd, *J* = 6.0, 4.0 Hz, 2H), 3.23 (br s, 4H), 2.27 (d, *J* = 4.0 Hz, 2H), 2.17 (septet, *J* = 6.0 Hz, 1H), 1.00 (d, *J* = 6.0 Hz, 6H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 173.7, 147.8, 133.4 (q, *J* = 31.5 Hz), 132.1, 123.9, 123.6 (q, *J* = 270.8 Hz), 117.1 (q, *J* = 3.8 Hz), 113.4 (q, *J* = 3.8 Hz), 47.8, 42.6, 41.3, 27.0, 22.9. LC-MS (*m/z*) calcd for C<sub>16</sub>H<sub>21</sub>F<sub>3</sub>N<sub>2</sub>O [M+H<sup>+</sup>], 315.2; found, 315.2. HPLC: purity<sub>254</sub> > 99%.

**3,3-Dimethyl-1-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)butan-1-one (2.9)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and 3,3-dimethylbutanoyl chloride by general procedure B in 73% yield as a pale-orange oil. *R*<sub>f</sub> 0.50 (heptane/

EtOAc 1:1). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.39–7.35 (m, 1H), 7.14–7.11 (m, 2H), 7.08–7.06 (m, 1H), 3.81 (dd, *J* = 8.0, 4.0 Hz, 2H), 3.69 (dd, *J* = 8.0, 4.0 Hz, 2H), 3.22 (dd, *J* = 8.0, 4.0 Hz, 2H), 3.21 (dd, *J* = 8.0, 4.0 Hz, 2H), 2.31 (s, 2H), 1.08 (s, 9H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 171.9, 147.3, 134.2 (q, *J* = 31.5 Hz), 132.3, 126.2, 124.7 (q, *J* = 270.8 Hz), 123.4 (q, *J* = 3.8 Hz), 117.4 (q, *J* = 3.8 Hz), 54.5, 46.1, 33.0, 31.5, 21.1. LC-MS (*m/z*) calcd for C<sub>17</sub>H<sub>23</sub>F<sub>3</sub>N<sub>2</sub>O [M+H<sup>+</sup>], 329.2; found, 329.2. HPLC: purity<sub>254</sub> > 96%.

**4-Methyl-1-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)pentan-1-one (2.10)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and 4-methylvaleryl chloride by general procedure B in 87% yield as a clear oil. *R*<sub>f</sub> 0.35 (heptane/EtOAc 1:1). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 7.37 (t, *J* = 6.0 Hz, 1H), 7.14–7.07 (m, 3H), 3.79 (dd, *J* = 8.0, 4.0 Hz, 2H), 3.65 (dd, *J* = 8.0, 4.0 Hz, 2H), 3.24 (dd, *J* = 8.0, 4.0 Hz, 2H), 3.22 (dd, *J* = 8.0, 4.0 Hz, 2H), 2.35 (dd, *J* = 6.0, 6.0 Hz, 2H), 1.63–1.52 (m, 3H), 0.93 (d, *J* = 3.0 Hz, 6H). <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 172.1, 151.1, 131.6 (q, *J* = 31.5 Hz), 129.7, 124.2 (q, *J* = 271.0 Hz), 119.3, 116.6 (q, *J* = 4.0 Hz), 112.7 (q, *J* = 4.0 Hz), 49.2, 49.0, 45.4, 41.3, 34.2, 31.3, 27.9, 22.7, 22.4. LC-MS (*m/z*) calcd for C<sub>17</sub>H<sub>23</sub>F<sub>3</sub>N<sub>2</sub>O [M+H<sup>+</sup>], 329.2; found, 329.2. HPLC: purity<sub>254</sub> > 99%.

**Phenyl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.11)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and benzoic acid by general procedure A in 58% yield as a pale-yellow oil. *R*<sub>f</sub> 0.33 (heptane/EtOAc 2:3). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.47–7.39 (m, 6H), 7.26–7.20 (m, 2H), 7.10 (d, *J* = 7.5 Hz, 1H), 3.75 (br s, 2H), 3.48 (br s, 2H), 3.28 (br s, 4H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 170.8, 151.5, 135.8, 132.1 (q, *J* = 31.5 Hz), 130.3 (d, *J* = 1.5 Hz), 130.1 (d, *J* = 1.5 Hz), 129.0, 127.5, 124.6 (q, *J* = 267.0 Hz), 119.9, 117.2, 113.3, 49.7, 47.8. LC-MS (*m/z*) calcd for C<sub>18</sub>H<sub>17</sub>F<sub>3</sub>N<sub>2</sub>O [M+H<sup>+</sup>], 335.1; found, 335.1. Anal. calcd for C<sub>18</sub>H<sub>17</sub>F<sub>3</sub>N<sub>2</sub>O × 1HCl: C 58.30, H 4.89, N 7.55 found C 58.20, H 4.78, N 7.43.

**Benzo[d][1,3]dioxol-5-yl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.12)**

Obtained from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and benzo[d][1,3]dioxole-5-carboxylic acid by general procedure A in 60% yield as a white solid. *R*<sub>f</sub> 0.18 (heptane/EtOAc 3:1). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.36 (t, *J* = 9.0 Hz, 1H), 7.13–7.05 (m, 3H), 6.97–6.82 (m, 3H), 6.00 (s, 2H), 3.77 (br s, 4H), 3.24 (br s, 4H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>) δ 169.9, 151.0, 149.0, 147.7, 131.5 (q, *J* = 31.5 Hz), 129.7, 128.9, 124.1 (q, *J* = 270.8 Hz), 121.7, 119.4, 116.8 (q, *J* = 3.8 Hz), 112.8 (q, *J* = 3.8 Hz), 108.2, 108.1, 101.5, 49.2, 42.8; mp 87–89°C. LC-MS (*m/z*)



calcd for  $C_{19}H_{17}F_3N_2O_3$  [ $M+H^+$ ], 379.1; found, 379.1. Anal. calcd for  $C_{19}H_{17}F_3N_2O_3$ : C 60.32, H 4.53, N 7.40 found C 59.95, H 4.40, N 7.02.

**(3-Phenoxyphenyl)(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.13)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (**4**) and 3-phenoxybenzoic acid by general procedure **A** in 62 % yield as a pale-yellow oil.  $R_f$  0.20 (heptane/EtOAc 3:1).  $^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.40–7.22 (m, 5H), 7.16–6.99 (m, 8H), 3.90 (br s, 2H), 3.60 (br s, 2H), 3.25 (br s, 2H), 3.16 (br s, 2H).  $^{13}C$  NMR (75 MHz,  $CDCl_3$ )  $\delta$  170.1, 158.1, 156.7, 151.4, 137.4, 132.0 (q,  $J = 31.5$  Hz), 130.6, 130.4, 130.1, 124.6 (q,  $J = 270.8$  Hz), 124.4, 121.9, 120.2, 119.9, 119.6, 117.3 (q,  $J = 3.8$  Hz), 113.3 (q,  $J = 3.8$  Hz), 49.7, 42.9. LC-MS ( $m/z$ ) calcd for  $C_{24}H_{21}F_3N_2O_2$  [ $M+H^+$ ], 427.1; found, 427.1. Anal. calcd for  $C_{24}H_{21}F_3N_2O_2 \times 1HCl$ : C 62.27, H 4.79, N 6.05 found C 63.52, H 4.64, N 5.90.

**1-Phenylcyclopentyl-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.14)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (**4**) and 1-phenylcyclopentanecarboxylic acid by general procedure **A** in 92 % yield as an off-white solid.  $R_f$  0.37 (heptane/EtOAc 3:1).  $^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.40–7.17 (m, 8H), 7.05 (d,  $J = 6.0$  Hz, 1H), 3.78 (br s, 4H), 3.25 (br s, 2H), 3.13 (br s, 2H), 2.70–2.41 (m, 4H), 2.06–1.87 (m, 4H).  $^{13}C$  NMR (75 MHz,  $CDCl_3$ )  $\delta$  175.2, 151.4, 145.8, 143.2, 131.9 (q,  $J = 31.5$  Hz), 130.0, 128.7, 127.5, 126.8, 124.6 (q,  $J = 270.8$  Hz), 119.4 (q,  $J = 1.5$  Hz), 116.8 (q,  $J = 3.8$  Hz), 112.9 (q,  $J = 3.8$  Hz), 59.3, 58.9, 48.7, 38.8, 36.4, 25.7, 24.0; mp 77–79°C. LC-MS ( $m/z$ ) calcd for  $C_{23}H_{25}F_3N_2O$  [ $M+H^+$ ], 403.2; found, 403.2. Anal. calcd for  $C_{23}H_{25}F_3N_2O$ : C 68.64, H 6.26, N 6.96 found C 70.90, H 6.37, N 6.73.

**Naphthalen-1-yl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.15)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (**4**) and 1-naphthoic acid by general procedure **A** in 83% yield as a colorless oil.  $R_f$  0.19 (heptane/EtOAc 3:1).  $^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.90–7.83 (m, 3H), 7.55–7.32 (m, 5H), 7.12–7.02 (m, 3H), 4.21–4.01 (m, 2H), 3.42–3.35 (m, 4H), 3.11–2.99 (m, 2H).  $^{13}C$  NMR (75 MHz,  $CDCl_3$ )  $\delta$  169.9, 151.4, 134.1, 133.9, 132.0 (q,  $J = 31.5$  Hz), 130.1, 129.8, 128.9, 127.6, 127.0, 125.6, 125.1, 124.5 (q,  $J = 270.8$  Hz), 124.3, 119.4 (q,  $J = 1.5$  Hz), 117.3 (q,  $J = 3.8$  Hz), 113.3 (q,  $J = 3.8$  Hz), 50.1, 49.7, 47.3, 42.0. LC-MS ( $m/z$ ) calcd for  $C_{22}H_{19}F_3N_2O$  [ $M+H^+$ ], 385.1; found, 385.1. Anal. calcd for  $C_{22}H_{19}F_3N_2O \times 1HCl$ : C 62.79, H 4.79, N 6.66 found C 64.82, H 4.72, N 6.26.

**Thiophen-2-yl(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.16)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (**4**) and thiophene-2-carboxylic acid by general procedure **A** in 69% yield as a pale-yellow oil.  $R_f$  0.68 (100% EtOAc).  $^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.46 (dd,  $J = 6.0, 3.0$  Hz, 1H), 7.39–7.32 (m, 2H), 7.13–7.04 (m, 4H), 3.92 (dd,  $J = 6.0, 3.0$  Hz, 4H), 3.28 (dd,  $J = 6.0, 3.0$  Hz, 4H).  $^{13}C$  NMR (75 MHz,  $CDCl_3$ )  $\delta$  164.1, 151.4, 137.1, 131.9 (q,  $J = 31.5$  Hz), 130.1, 129.5, 129.3, 127.2, 124.6 (q,  $J = 270.8$  Hz), 119.7 (q,  $J = 1.5$  Hz), 117.2 (q,  $J = 3.8$  Hz), 113.1 (q,  $J = 3.8$  Hz), 49.5, 45.7. LC-MS ( $m/z$ ) calcd for  $C_{16}H_{15}F_3N_2OS$  [ $M+H^+$ ], 341.0; found, 341.0. Anal. calcd for  $C_{16}H_{15}F_3N_2OS \times 1HCl$ : C 51.00, H 4.28, N 7.43 found C 51.41, H 4.28, N 7.33.

**1-(4-(3-(Trifluoromethyl)phenyl)piperazin-1-yl)-2-phenylethanone (2.17)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (**4**) and 2-phenylacetyl chloride by general procedure **B** in 77% yield as a clear oil.  $R_f$  0.28 (heptane/EtOAc 1:1).  $^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.34–7.20 (m, 6H), 7.09–6.96 (m, 3H), 3.79 (dd,  $J = 6.0, 3.0$  Hz, 2H), 3.78 (s, 2H), 3.58 (dd,  $J = 6.0, 3.0$  Hz, 2H), 3.16 (dd,  $J = 6.0, 3.0$  Hz, 2H), 2.99 (dd,  $J = 6.0, 3.0$  Hz, 2H).  $^{13}C$  NMR (75 MHz,  $CDCl_3$ )  $\delta$  169.8, 151.3, 137.8, 131.3 (q,  $J = 31.5$  Hz), 129.4, 129.1, 128.3, 127.1, 124.3 (q,  $J = 270.8$  Hz), 118.6 (d,  $J = 0.8$  Hz), 115.6 (q,  $J = 3.8$  Hz), 112.0 (q,  $J = 3.8$  Hz), 62.9, 52.8, 48.6. LC-MS ( $m/z$ ) calcd for  $C_{19}H_{19}F_3N_2O$  [ $M+H^+$ ], 349.1; found, 349.1. Anal. Calcd for  $C_{19}H_{19}F_3N_2O \times 1HCl$ : C 60.20, H 5.16, N 7.12 found C 59.30, H 5.24, N 7.28.

**4-Aminophenyl-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (2.18)**

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (**4**) (103 mg, 0.37 mmol) and 4-((*tert*-butoxycarbonyl)amino)benzoic acid in accordance with general procedure **A** to provide *N*-Boc-**2.18** in 48% yield as a brown oil.  $R_f$  0.44 (heptane/EtOAc 1:1).  $^1H$  NMR (400 MHz, MeOD)  $\delta$  7.37–7.27 (m, 5H), 7.07–6.99 (m, 3H), 3.81–3.61 (m, 4H), 3.27–3.07 (m, 4H), 1.46 (s, 9H).  $^{13}C$  NMR (100 MHz, MeOD)  $\delta$  170.3, 152.5, 151.1, 140.1, 131.8, 131.5, 129.7, 128.5, 127.5, 125.6, 119.4, 118.1, 116.8, 114.1, 112.9 (d,  $J = 4.0$  Hz), 81.1, 49.2, 31.9, 29.0, 28.3, 22.7. LC-MS ( $m/z$ ) calcd for  $C_{23}H_{26}F_3N_3O_3$  [ $M+H^+$ ], 450.1, found, 450.1. The intermediate product *N*-Boc-**2.18** was dissolved in DCM (2 mL) and TFA (2 mL) and stirred for 1h at rt. The reaction mixture was evaporated and the title compound **2.18** was obtained in 69% yield as TFA-salt.  $^1H$  NMR (400 MHz, MeOD)  $\delta$  7.43–7.39 (m, 1H), 7.35–7.32 (m, 2H), 7.23–7.20 (m, 2H), 7.12–7.10 (m, 1H), 6.90–6.87 (m, 2H), 3.79 (s, 4H), 3.30–3.26 (m, 4H).  $^{13}C$  NMR (100 MHz, MeOD)  $\delta$  173.1, 152.1, 148.3, 132.8, 132.5 (q,  $J = 8.0$  Hz), 131.0, 130.4, 127.2, 126.7, 124.5, 120.8, 118.6, 117.2

(q,  $J = 1.0$  Hz), 117.0, 115.0, 113.6 (q,  $J = 1.0$  Hz), 50.1, 49.7. LC-MS ( $m/z$ ) calcd for  $C_{18}H_{18}F_3N_3O$  [ $M+H^+$ ], 350.1; found, 350.1. HPLC: purity<sub>254</sub> >97%.

### 3-Aminophenyl-4-(3-(trifluoromethyl)phenyl)piperazin-1-yl) methanone (2.19)

Prepared from 3-trifluoromethylphenylpiperazine (4) (101 mg, 0.37 mmol) and 3-((*tert*-butoxycarbonyl)amino) benzoic acid by general procedure A in 85% yield as a reddish oil.  $R_f$  0.42 (heptane/EtOAc 1:1).  $^1H$  NMR (400 MHz, MeOD)  $\delta$  7.45 (s, 1H), 7.36–7.23 (p,  $J = 8.0$  Hz, 3H), 7.06–6.98 (m, 4 H), 6.78 (s, 1H), 3.95–3.75 (m, 2H), 3.66–3.46 (m, 2H), 3.27–3.01 (m, 4H), 1.45 (s, 9H).  $^{13}C$  NMR (100 MHz, MeOD)  $\delta$  172.4, 155.2, 152.9, 141.1 (q,  $J = 2.0$  Hz), 137.2, 132.6 (q,  $J = 4.0$  Hz), 131.0, 130.3, 129.8, 127.2, 124.8, 122.0, 120.9 (q,  $J = 2.0$  Hz), 118.0, 117.3, 113.7, 81.2, 79.5, 54.8, 30.7, 28.7, 24.3, 23.8. LC-MS ( $m/z$ ) calcd for  $C_{23}H_{26}F_3N_3O_3$  [ $M+H^+$ ], 450.1, found, 450.1. The intermediate product *N*-Boc-2.19 was dissolved in DCM (2 mL) and TFA (2 mL) and stirred for 1h at rt. The reaction mixture was evaporated to afford the title compound in 99% yield as TFA-salt.  $^1H$  NMR (400 MHz, MeOD)  $\delta$  7.67–7.62 (m, 2H), 7.57–7.49 (m, 3H), 7.44–7.40 (m, 1H), 7.24–7.22 (m, 2H), 7.14–7.12 (m, 1H), 4.01–3.80 (m, 2H), 3.74–3.50 (m, 2H), 3.45–3.18 (m, 4H).  $^{13}C$  NMR (100 MHz, MeOD)  $\delta$  170.8, 159.0 (q,  $J = 10.5$  Hz), 152.7, 138.6, 134.6, 134.24, 133.5, 132.5 (q,  $J = 8.0$  Hz), 131.7, 131.5, 131.01, 130.6, 128.5, 127.5, 127.2, 125.0, 124.9, 124.5, 122.3, 120.9 (d,  $J = 1.0$  Hz), 117.4 (q,  $J = 1.0$  Hz), 114.7, 113.7 (q,  $J = 1.0$  Hz), 54.5, 54.2. LC-MS ( $m/z$ ) calcd for  $C_{18}H_{18}F_3N_3O$  [ $M+H^+$ ], 350.1; found, 350.1. HPLC: purity<sub>254</sub> >96%.

### 2-Amino-1-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl) ethanone (2.20)

Prepared from (4) (104 mg, 0.37 mmol) and *N*-Boc-glycine-OH (105 mg, 0.36 mmol) by general procedure A to provide *N*-Boc-2.20 in 86% yield as a colorless oil.  $R_f$  0.25 (heptane/EtOAc 1:1).  $^1H$  NMR (400 MHz, MeOD)  $\delta$  7.41 (t,  $J = 8.0$  Hz, 1H), 7.23–7.20 (m, 2H), 7.11 (d,  $J = 8.0$  Hz, 1H), 3.97 (s, 2H), 3.75 (s, 2H), 3.66 (s, 2H), 3.28–3.23 (m, 4H), 1.45 (s, 9H). The intermediate *N*-Boc-2.20 was dissolved in DCM (2 mL) and TFA (2 mL) and stirred for 30 min at rt. The reaction mixture was evaporated to afford the title compound as the TFA-salt in quantitative yield.  $^1H$  NMR (400 MHz, MeOD)  $\delta$  7.45–7.40 (m, 1H), 7.24–7.21 (m, 2H), 7.14–7.12 (m, 1H), 4.00 (s, 2H), 3.80 (t,  $J = 4.0$  Hz, 2H), 3.62 (t,  $J = 4.0$  Hz, 2H), 3.30–3.26 (m, 2H).  $^{13}C$  NMR (100 MHz, MeOD)  $\delta$  165.7, 159.9 (q,  $J = 41.0$  Hz), 152.7, 132.40 (t, 32.0 Hz), 131.0, 127.2, 124.5, 120.9, 117.5, 114.7, 113.6 (q,  $J = 4.0$  Hz), 49.9, 49.7, 45.5, 43.1, 41.0, 27.7. LC-MS ( $m/z$ ) calcd for  $C_{13}H_{16}F_3N_3O$  [ $M+H^+$ ], 288.1; found, 288.1. HPLC: purity<sub>254</sub> >99%.

### 2-Amino-1-(4-(3-(trifluoromethyl)phenyl)piperazin-1-yl) ethanone (2.21)

Prepared from (16) (103 mg, 0.37 mmol) and *N*-Boc-glycine-OH (69 mg, 0.36 mmol) by general procedure A to provide *N*-Boc-2.21 in 51% yield as a colorless oil.  $R_f$  0.18 (heptane/EtOAc 1:1). The intermediate *N*-Boc-2.20 was dissolved in DCM (2 mL) and TFA (2 mL) and stirred for 45 min at rt. The reaction mixture was evaporated to afford the title compound as the TFA-salt in quantitative yield.  $^1H$  NMR (400 MHz, MeOD)  $\delta$  7.44–7.40 (m, 1H), 7.24–7.20 (m, 2H), 7.12 (dt,  $J = 8.0$  Hz, 1H), 3.78 (t,  $J = 4.0$  Hz, 2H), 3.68 (t,  $J = 4.0$  Hz, 2H), 3.26–3.21 (m, 6H), 2.83 (t,  $J = 4.0$  Hz, 2H).  $^{13}C$  NMR (100 MHz, MeOD)  $\delta$  170.4, 152.8, 132.7, 132.4, 131.0, 127.2, 120.8, 117.5, 117.3 (q,  $J = 4.0$  Hz), 113.5 (q,  $J = 4.0$  Hz), 54.4, 50.0, 46.16, 42.65. LC-MS ( $m/z$ ) calcd for  $C_{14}H_{18}F_3N_3O$  [ $M+H^+$ ], 302.1; found, 302.1. HPLC: purity<sub>254</sub> >98%.

### Benzyl 4-(3-(trifluoromethyl)phenyl)piperazine-1-carboxylate (2.32)

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and benzyl carbonochloridate by general procedure B in 73% yield as a clear oil.  $R_f$  0.43 (heptane/EtOAc 3:1).  $^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.36–7.29 (m, 6H), 7.10–7.01 (m, 3H), 5.15 (s, 2H), 3.66 (dd,  $J = 6.0, 3.0$  Hz, 4H), 3.17 (br s, 4H).  $^{13}C$  NMR (75 MHz,  $CDCl_3$ )  $\delta$  155.1, 151.2, 136.5, 131.3 (q,  $J = 31.5$  Hz), 129.6, 128.5, 128.1, 127.9, 124.2 (q,  $J = 270.8$  Hz), 119.4 (q,  $J = 1.5$  Hz), 116.5 (q,  $J = 3.8$  Hz), 112.4 (q,  $J = 3.8$  Hz), 67.3, 48.9, 43.5. LC-MS ( $m/z$ ) calcd for  $C_{19}H_{19}F_3N_2O_2$  [ $M+H^+$ ], 365.1; found, 365.1. Anal. Calcd for  $C_{19}H_{19}F_3N_2O_2 \times 1HCl$ : C 57.35, H 5.03, N 6.91 found C 56.93, H 5.03, N 6.99.

### 1-(Phenylsulfonyl)-4-(3-(trifluoromethyl)phenyl)piperazine (2.33)

Prepared from 1-(3-(trifluoromethyl)phenyl)piperazine (4) and benzenesulfonyl chloride by general procedure B in 71% yield as a white solid.  $R_f$  0.39 (heptane/EtOAc 3:1).  $^1H$  NMR (300 MHz,  $CDCl_3$ )  $\delta$  7.80–7.76 (m, 2H), 7.65–7.51 (m, 3H), 7.32 (t,  $J = 7.5$  Hz, 1H), 7.11–6.97 (m, 3H), 3.30–3.26 (m, 4H), 3.19–3.16 (m, 4H).  $^{13}C$  NMR (75 MHz,  $CDCl_3$ )  $\delta$  151.1, 135.6, 133.5, 131.9 (q,  $J = 31.5$  Hz), 130.2, 129.6, 128.2, 124.5 (q,  $J = 270.8$  Hz), 119.9 (q,  $J = 1.5$  Hz), 117.4 (q,  $J = 3.8$  Hz), 113.5 (q,  $J = 3.8$  Hz), 49.1, 46.4; mp 109–111°C (decomposed). LC-MS ( $m/z$ ) calcd for  $C_{17}H_{17}F_3N_2O_2S$  [ $M+H^+$ ], 371.1; found, 371.1. Anal. Calcd for  $C_{17}H_{17}F_3N_2O_2S$ : C 55.13, H 4.63, N 7.56 found C 54.60, H 4.44, N 7.41.

### (±)-endo-exo-Bicyclo[2.2.1]heptan-2-ylmethyl)-4-(3-(trifluoromethyl)phenyl)piperazine (2.34)

A solution of (±)-endo-exo-bicyclo[2.2.1]heptan-2-yl (4-(3-(trifluoromethyl)phenyl)piperazin-1-yl)methanone ((±)-endo-exo 2.3) (116.5 mg, 0.30 mmol) in dry THF

(3 mL) was added dropwise to a solution of  $\text{LiAlH}_4$  (13 mg, 0.33 mmol) in THF (5 mL) at  $0^\circ\text{C}$  under a  $\text{N}_2$  atmosphere. The reaction mixture was stirred for 3 days at rt, quenched with 2N NaOH (2 mL) and extracted with dichloromethane ( $3 \times 25$  mL). The combined organic phases were washed with  $\text{H}_2\text{O}$  (20 mL) and brine (20 mL). The organic phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$ . After concentration *in vacuo*, the crude product was purified by column chromatography on silica gel to afford the titled compound as a pale-yellow oil (72 mg, 0.21 mmol, 71% yield):  $R_f$  0.42 (heptane/EtOAc 2:1).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.37 (t,  $J = 8.0$  Hz, 1H), 7.13–7.00 (m, 3H), 3.91–3.66 (m, 4H), 3.25–3.16 (m, 4H), 2.97–2.41 (m, 2H), 2.32–2.28 (m, 1H), 1.98–1.92 (m, 1H), 1.62–1.19 (m, 9H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  151.2, 130.1 (q,  $J = 31.5$  Hz), 130.2, 124.5 (q,  $J = 270.7$  Hz), 119.3 (d,  $J = 3.8$  Hz), 117.3 (q,  $J = 3.8$  Hz), 113.2 (q,  $J = 3.8$  Hz), 49.0, 44.3, 40.9, 40.4, 37.2, 36.7, 36.0, 34.9, 32.2, 29.5, 28.9, 24.5. LC-MS ( $m/z$ ) calcd for  $\text{C}_{19}\text{H}_{25}\text{F}_3\text{N}_2$  [ $\text{M}+\text{H}^+$ ], 339.2; found, 339.2. HPLC: purity<sub>254</sub> > 99%.

#### 1-Benzyl-4-(3-(trifluoromethyl)phenyl)piperazine (2.35)

BnBr (54  $\mu\text{L}$ , 0.46 mmol) was added dropwise to a solution of 1-(3-(trifluoromethyl)phenyl)piperazine (**4**) (100 mg, 0.43 mmol) and  $\text{Et}_3\text{N}$  (126  $\mu\text{L}$ , 0.46 mmol) in dichloromethane (5 mL) at rt under a  $\text{N}_2$  atmosphere. The reaction mixture was stirred for 24 hours, quenched with saturated  $\text{NH}_4\text{Cl}$  (5 mL) and extracted with dichloromethane ( $3 \times 20$  mL). The combined organic phases were washed with  $\text{H}_2\text{O}$  (20 mL) and brine (20 mL). The organic phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$ . After concentration *in vacuo*, the crude product was purified by column chromatography on silica gel to afford the titled compound as a clear oil (79 mg, 0.25 mmol, 57% yield):  $R_f$  0.41 (heptane/EtOAc 2:1).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.34–7.19 (m, 6H), 7.08–6.99 (m, 3H), 3.55 (br s, 2H), 3.21 (dd,  $J = 6.0$ , 6.0 Hz, 4H), 2.59 (dd,  $J = 6.0$ , 6.0 Hz, 4H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  151.3, 135.2, 131.7 (q,  $J = 31.5$  Hz), 130.1, 129.3, 128.9, 127.4, 124.6 (q,  $J = 270.8$  Hz), 119.6 (d,  $J = 3.8$  Hz), 117.0 (q,  $J = 3.8$  Hz), 113.0 (q,  $J = 3.8$  Hz), 49.3, 49.1, 46.2, 41.9, 41.5. LC-MS ( $m/z$ ) calcd for  $\text{C}_{18}\text{H}_{19}\text{F}_3\text{N}_2$  [ $\text{M}+\text{H}^+$ ], 321.2; found, 321.2. Anal. Calcd for  $\text{C}_{18}\text{H}_{19}\text{F}_3\text{N}_2 \times \text{HCl}$ : C 60.59, H 5.65, N 7.85 found C 55.74, H 5.23, N 7.12.

#### Bicyclo[2.2.1]heptan-2-yl(4-phenylpiperazin-1-yl)methanone (3.1)

Prepared in accordance with *general procedure A*. The crude product was purified by flash chromatography to afford **3.1** as a clear oil (240 mg, 43%).  $R_f$  0.20 (heptane/EtOAc 3:1).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.31–7.25 (m, 2H), 6.95–6.85 (m, 3H), 3.99–3.61 (m, 4H), 3.30–3.03 (m, 4H), 2.98–2.91 (m, 0.5H), 2.44–2.39 (m, 1.5H), 2.34–2.26 (m, 1H), 1.99–1.90 (m, 1H), 1.64–1.16 (m, 7H).

$^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  173.9, 172.2, 151.0, 129.2, 120.5, 120.5, 116.6, 50.2, 49.8, 49.7, 49.6, 45.4, 45.3, 44.2, 43.7, 41.9, 41.7, 40.8, 40.6, 40.4, 37.2, 36.7, 36.0, 34.9, 32.2, 29.5, 28.9, 28.9, 24.5. LC-MS ( $m/z$ ) calcd for  $\text{C}_{18}\text{H}_{24}\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 285.1; found, 285.1. HPLC: purity<sub>254</sub> > 95%.

#### ( $\pm$ )-endo-exo-Bicyclo[2.2.1]heptan-2-yl(4-(4-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (3.2)

$\text{Pd}(\text{OAc})_2$  (1.0 mg, 4.6  $\mu\text{mol}$ ) and  $\text{P}^t\text{Bu}_3$  (1.0 M in toluene, 16  $\mu\text{L}$ , 0.016 mmol) were added to a solution of 4-(bicyclo[2.2.1]heptane-2-carbonyl)piperazin-1-ium chloride ( $\pm$ )-endo-exo-7 (101 mg, 0.41 mmol), 1-bromo-4-(trifluoromethyl)benzene (63  $\mu\text{L}$ , 0.45 mmol) and  $\text{NaO}^t\text{Bu}$  (86 mg, 0.90 mmol) in dry *o*-xylene (1.3 mL) at rt under a  $\text{N}_2$  atmosphere. The reaction mixture was stirred at  $120^\circ\text{C}$  for 24 hours, quenched with  $\text{H}_2\text{O}$  (10 mL) and extracted with EtOAc ( $3 \times 20$  mL). The combined organic phases were washed with  $\text{H}_2\text{O}$  (20 mL) and brine (20 mL). The organic phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$ . After concentration *in vacuo*, the crude product was purified by column chromatography on silica gel to afford the titled compound as a white solid (80 mg, 0.23 mmol, 56%):  $R_f$  0.25 (heptane/EtOAc 3:1).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.42 (d,  $J = 8.6$  Hz, 2H), 6.86 (d,  $J = 8.8$  Hz, 2H), 3.90–3.80 (m, 0.5H), 3.75–3.62 (m, 4H), 3.31–3.08 (m, 4H), 2.91–2.80 (m, 0.5H), 2.40–2.30 (m, 1H), 2.28–2.19 (m, 1H), 1.91–1.82 (m, 1H), 1.60–1.10 (m, 7H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  174.0, 172.3, 153.0, 126.6, 126.5, 126.4, 126.4, 125.9, 123.2, 121.4, 121.1, 114.9, 48.8, 48.4, 48.3, 48.2, 45.1, 44.9, 44.2, 43.7, 41.6, 41.4, 40.8, 40.3, 37.1, 36.7, 36.0, 34.9, 32.2, 29.5, 28.9, 28.9, 24.5; mp:  $126\text{--}127^\circ\text{C}$  (decomposed). LC-MS ( $m/z$ ) calcd for  $\text{C}_{19}\text{H}_{23}\text{F}_3\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 353.2; found, 353.2. HPLC: purity<sub>254</sub> > 95%.

#### ( $\pm$ )-endo-exo-Bicyclo[2.2.1]heptan-2-yl(4-(2-(trifluoromethyl)phenyl)piperazin-1-yl)methanone (( $\pm$ )-endo-exo-3.5)

Prepared in accordance with *general procedure A*. The crude product was purified twice by flash chromatography to afford ( $\pm$ )-endo-exo-**3.5** as a clear oil (48 mg, 14%).  $R_f$  0.32 (heptane/EtOAc 3:1).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.64 (d,  $J = 7.86$  Hz, 1H), 7.52 (t,  $J = 7.69$  Hz, 1H), 7.31 (d,  $J = 8.02$  Hz, 1H), 7.28–7.22 (m, 1H), 3.72 (s, 4H), 2.99–2.85 (m, 4.5H), 2.45–2.36 (m, 1.5H), 2.33–2.25 (m, 1H), 2.00–1.90 (m, 1H), 1.65–1.15 (m, 7H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  173.9, 172.3, 151.8, 132.8, 127.6, 127.3, 127.2, 125.2, 124.0, 122.6, 54.1, 53.8, 53.3, 53.2, 46.0, 45.9, 44.2, 43.7, 42.4, 42.2, 40.8, 40.6, 40.3, 37.2, 36.7, 36.0, 34.9, 32.2, 29.4, 28.9, 28.9, 24.5. LC-MS ( $m/z$ ) calcd for  $\text{C}_{19}\text{H}_{23}\text{F}_3\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 353.1; found, 353.1. HPLC: purity<sub>254</sub> > 95%.

#### ( $\pm$ )-endo-exo-Bicyclo[2.2.1]heptan-2-yl(4-(3-chlorophenyl)piperazin-1-yl)methanone (( $\pm$ )-endo-exo-3.7)

Prepared in accordance with *general procedure A*. The crude product was purified by flash chromatography to



afford **3.7** as a clear oil (90 mg, 46%).  $R_f$  0.31 (heptane/EtOAc 3:1).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.17 (t,  $J = 8.09$  Hz, 1H), 6.87 (t,  $J = 2.04$  Hz, 1H), 6.85 (d,  $J = 7.84$  Hz, 1H), 6.78 (dd,  $J = 8.35, 2.15$  Hz, 1H), 3.95–3.60 (m, 4H), 3.25–3.05 (m, 4H), 2.97–2.90 (m, 0.5H), 2.45–2.37 (m, 1.5H), 2.33–2.25 (m, 1H), 1.98–1.88 (m, 1H), 1.64–1.06 (m, 7H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  173.8, 172.1, 151.9, 134.9, 130.0, 119.9, 116.2, 114.3, 49.4, 49.1, 48.9, 45.1, 44.9, 44.9, 43.6, 41.5, 41.3, 40.7, 40.5, 40.2, 37.0, 36.6, 35.9, 34.8, 32.0, 29.3, 28.7, 24.4. LC-MS ( $m/z$ ) calcd for  $\text{C}_{18}\text{H}_{23}\text{ClN}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 319.1; found, 319.1. HPLC: purity<sub>254</sub> > 97%.

**( $\pm$ )-endo-exo-Bicyclo[2.2.1]heptan-2-yl(4-(3-hydroxyphenyl)piperazin-1-yl)methanone (( $\pm$ )-endo-exo-3.8)**

Prepared from ( $\pm$ )-endo-exo-bicyclo[2.2.1]heptane-2-carboxylic acid (( $\pm$ )-endo-exo-6) and 3-(piperazin-1-yl)phenol by general procedure A in 40% yield as a white solid.  $R_f$  0.45 (heptane/EtOAc 2:1).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.13 (t,  $J = 8.0$  Hz, 1H), 6.52–6.36 (m, 3H), 3.93–3.64 (m, 4H), 3.22–2.92 (m, 4H), 2.45–2.41 (m, 1H), 2.31–2.28 (m, 1H), 1.97–1.88 (m, 1H), 1.63–1.18 (m, 8H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  173.5, 157.4, 152.3, 130.1, 108.7, 107.7, 103.6, 50.2, 49.8, 49.1, 45.3, 44.3, 43.7, 41.8, 40.6, 40.5, 37.2, 36.8, 36.0, 35.1, 32.2, 29.5, 28.9, 28.9, 24.5; mp: 187–189°C (decomposed). LC-MS ( $m/z$ ) calcd for  $\text{C}_{18}\text{H}_{24}\text{N}_2\text{O}_2$  [ $\text{M}+\text{H}^+$ ], 301.2; found, 301.2. HPLC: purity<sub>254</sub> > 98%.

**3-(4-(bicyclo[2.2.1]heptane-2-carbonyl)piperazin-1-yl)benzonitrile (3.9)**

Prepared in accordance with general procedure A. The crude product was purified by flash chromatography to afford **3.9** as a clear oil (63 mg, 63%).  $R_f$  0.15 (heptane/EtOAc 3:1).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.37–7.31 (m, 1H), 7.15–7.10 (m, 3H), 3.98–3.62 (m, 4H), 3.30–3.10 (m, 4H), 2.98–2.90 (m, 0.5H), 2.46–2.36 (m, 1.5H), 2.34–2.26 (m, 1H), 1.98–1.88 (m, 1H), 1.64–1.10 (m, 7H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  173.9, 172.3, 151.0, 130.0, 123.3, 120.4, 119.1, 118.9, 113.2, 49.1, 48.8, 48.6, 45.0, 44.9, 44.2, 43.7, 41.5, 41.3, 40.8, 40.6, 40.3, 37.1, 36.7, 36.0, 34.9, 32.2, 29.4, 28.9, 28.8, 24.5. LC-MS ( $m/z$ ) calcd for  $\text{C}_{19}\text{H}_{23}\text{N}_3\text{O}$  [ $\text{M}+\text{H}^+$ ], 310.1; found, 310.1. HPLC: purity<sub>254</sub> > 99%.

**( $\pm$ )-endo-exo-Bicyclo[2.2.1]heptan-2-yl(4-(3-methoxyphenyl)piperazin-1-yl)methanone (( $\pm$ )-endo-exo-3.10)**

Prepared from ( $\pm$ )-endo-exo-bicyclo[2.2.1]heptane-2-carboxylic acid (( $\pm$ )-endo-exo-6) and 1-(3-methoxyphenyl)piperazine by general procedure A in 45% yield as a clear oil.  $R_f$  0.35 (heptane/EtOAc 2:1).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.19 (t,  $J = 8.0$  Hz, 1H), 6.55–6.44 (m, 3H), 3.92–3.61 (m, 4H), 3.48 (s, 3H), 3.24–2.92 (m, 4H), 2.44–2.40 (m, 1H), 2.31–2.27 (m, 1H), 1.97–1.89 (m, 1H), 1.61–1.17 (m, 8H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  173.2,

160.7, 152.4, 129.9, 109.3, 105.1, 103.1, 55.2, 50.8, 50.0, 49.6, 49.6, 49.4, 45.4, 45.3, 44.3, 43.8, 41.9, 41.7, 40.8, 40.6, 40.4, 37.2, 36.8, 36.0, 34.9, 32.2, 29.5, 29.0, 28.9, 27.0, 24.5. LC-MS ( $m/z$ ) calcd for  $\text{C}_{19}\text{H}_{26}\text{N}_2\text{O}_2$  [ $\text{M}+\text{H}^+$ ], 315.2; found, 315.2. HPLC: purity<sub>254</sub> > 98%.

**( $\pm$ )-endo-exo-Bicyclo[2.2.1]heptan-2-yl(4-(2,4-difluorophenyl)piperazin-1-yl)methanone (( $\pm$ )-endo-exo-3.11)**

Prepared from ( $\pm$ )-endo-exo-bicyclo[2.2.1]heptane-2-carboxylic acid (( $\pm$ )-endo-exo-6) and 1-(2,4-difluorophenyl)piperazine by general procedure A. After chromatography ( $R_f = 0.43$ , heptane/EtOAc 2:1). The title compound was isolated in 58% yield as a clear oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  6.92–6.77 (m, 3H), 3.93–3.64 (m, 4H), 3.07–2.90 (m, 4H), 2.44–2.27 (m, 2H), 1.98–1.90 (m, 1H), 1.62–1.17 (m, 8H).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  173.9, 172.3, 159.5 (d,  $J = 12.0$  Hz), 157.0 (d,  $J = 12.0$  Hz), 154.6 (d,  $J = 12.0$  Hz), 136.4 (d,  $J = 4.0$  Hz), 136.2 (d,  $J = 4.0$  Hz), 129.0, 128.4, 126.9, 119.9 (q,  $J = 4.0$  Hz), 110.9 (d,  $J = 2.0$  Hz), 110.7 (d,  $J = 2.0$  Hz), 105.1, 104.9 (d, 2.0 Hz), 104.6, 53.4, 51.8, 51.4, 51.0, 45.7, 45.6, 44.2, 43.2, 42.0, 41.8, 40.8, 40.6, 40.4, 37.2, 36.8, 36.0, 29.0, 28.9, 24.6. LC-MS ( $m/z$ ) calcd for  $\text{C}_{18}\text{H}_{22}\text{F}_2\text{N}_2\text{O}$  [ $\text{M}+\text{H}^+$ ], 321.2; found, 321.2. HPLC: purity<sub>254</sub> > 97%.

**1-(3-(Trifluoromethyl)phenyl)piperazine (4)**

Piperazine (5.74 g, 66.6 mmol), 3-bromobenzotrifluoride (5.0 g, 22.2 mmol),  $\text{Pd}(\text{OAc})_2$  (50 mg, 220  $\mu\text{mol}$ ),  $\text{P}(t\text{Bu})_3$  (216  $\mu\text{L}$ , 880  $\mu\text{mol}$ ) and  $\text{NaO}^t\text{Bu}$  (3 g, 31.1 mmol) were stirred in dry *o*-xylene (50 mL) at 120°C under a  $\text{N}_2$  atmosphere for 17 h.  $\text{H}_2\text{O}$  (25 mL) was added and the crude reaction was extracted with EtOAc (3  $\times$  50 mL) and the combined organic phases were washed with  $\text{H}_2\text{O}$  (30 mL) and brine (30 mL). The organic phase was dried over anhydrous  $\text{Na}_2\text{SO}_4$ . After concentration *in vacuo*, the crude product was purified by column chromatography on silica gel. This afforded the titled compound (3.91 g, 17.0 mmol, 77%) as a yellow oil:  $R_f$  0.23 ( $\text{Et}_2\text{O}/\text{MeCN}/\text{MeOH}/\text{Et}_3\text{N}$  10:1:1:0.5).  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  7.33 (t,  $J = 7.95$  Hz, 1H), 7.10–7.03 (m, 3H), 3.18 (dd,  $J = 9.0, 5.4$  Hz, 4H), 3.04 (dd,  $J = 6.9, 5.1$  Hz, 4H), 2.0 (s, 1H).  $^{13}\text{C}$  NMR (75 MHz,  $\text{CDCl}_3$ )  $\delta$  152.2, 131.9 (q,  $J = 30.8$  Hz), 129.9, 124.7 (q,  $J = 271.5$  Hz), 119.1 (q,  $J = 1.5$  Hz), 116.2 (q,  $J = 3.8$  Hz), 112.5 (q,  $J = 4.5$  Hz), 50.1, 46.3. LC-MS ( $m/z$ ) calcd for  $\text{C}_{11}\text{H}_{13}\text{F}_3\text{N}_2$  [ $\text{M}+\text{H}^+$ ], 231.1; found, 231.1. HPLC: purity<sub>254</sub> > 95%.

**( $\pm$ )-endo-exo-Bicyclo[2.2.1]heptane-2-carboxylic acid (( $\pm$ )-endo-exo-6)**

A solution of  $\text{KMnO}_4$  (53.2 g, 336.5 mmol) in  $\text{H}_2\text{O}$  (190 mL) was added dropwise to a solution of bicyclo[2.2.1]heptan-2-ylmethanol (17.0 g, 134.6 mmol) and  $\text{K}_2\text{CO}_3$  (7.4 g, 53.8 mmol) in  $\text{H}_2\text{O}$  (380 mL) at 0°C and stirred at room temperature for 24 hours. The crude

reaction was quenched with 4N HCl (pH  $\approx$  2) and extracted with EtOAc (3  $\times$  1000 mL). The combined organic phases were washed with H<sub>2</sub>O (500 mL), brine (250 mL) and dried over anhydrous MgSO<sub>4</sub>. After concentration *in vacuo*, the crude product was directly used for the next step without purification.

**( $\pm$ )-endo-exo-Bicyclo[2.2.1]heptan-2-yl(piperazin-1-yl) methanone (( $\pm$ )-endo-exo-7)**

DIPEA (2.8 g, 21.4 mmol) was added dropwise to a stirred solution of ( $\pm$ )-endo-exo-bicyclo[2.2.1]heptane-2-carboxylic acid (( $\pm$ )-endo-exo-7) (1 g, 7.1 mmol) and piperazine (1.85 g, 21.4 mmol) in dry DMF (60 mL) at 0°C under a N<sub>2</sub> atmosphere. A solution of TBTU (2.98 g, 9.27 mmol) in DMF (26 mL) was added to the reaction mixture and stirred at 0°C for 30 min. After 20 hours at rt, the reaction mixture was quenched with brine (30 mL) and extracted with dichloromethane (3  $\times$  100 mL). The combined organic phases were washed with H<sub>2</sub>O (3  $\times$  200 mL) and brine (100 mL). The organic phase was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>. After concentration *in vacuo*, the crude product was purified by column chromatography on silica gel (Et<sub>2</sub>O/MeOH/MeCN/Et<sub>3</sub>N 8:2:2:0.5). This afforded the title compound as a pale-yellow oil (0.73 mg, 3.5 mmol, 49% yield). The pure product was dissolved in dichloromethane (40 mL) and 4N HCl in dioxane (0.88 mL, 3.5 mmol) was added. The solution was evaporated to afford the corresponding HCl salt. <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  10.10 (br s, 2H), 4.10–3.80 (m, 4H), 3.22 (br s, 4H), 2.95–2.83 (m, 0.5H), 2.40–2.27 (m, 2.5H), 1.97–1.80 (m, 1H), 1.70–1.16 (m, 6H). <sup>13</sup>C NMR (75 MHz, CDCl<sub>3</sub>)  $\delta$  173.8, 172.2, 45.9, 45.4, 45.2, 45.0, 44.9, 44.8, 44.0, 43.5, 41.2, 41.1, 40.7, 40.4, 40.1, 37.0, 36.6, 35.8, 34.7, 32.0, 29.3, 28.7, 24.4, 8.7.

Analogues **2.22–2.31**, **2.36–2.40**, **3.3**, **3.4**, **3.6** and **3.12** were obtained as 5 mg portions pre-dissolved in DMSO from ChemBridge corporation and used directly in the [3H]-Asp-uptake assay.

**Abbreviations**

CNS: Central nervous system; EAAT: Excitatory amino acid transporter; SAR: Structure-activity-relationship.

**Competing interests**

All authors declare no financial competing interests.

**Authors' contributions**

THVH, molecular design, synthesis and manuscript preparation. CSD, molecular design, synthesis and manuscript preparation. BA, molecular design, pharmacological characterization and manuscript preparation. EM, synthesis. MF, synthesis. AAJ, molecular design, pharmacological characterization and manuscript preparation. LB, molecular design and manuscript preparation. All authors read and approved the final manuscript.

**Acknowledgements**

We would like to thank the Lundbeck Foundation, the Carlsberg Foundation, the Novo Nordisk Foundation, and the Danish Medical Research Council for financial support.

Received: 24 January 2013 Accepted: 28 February 2013

Published: 14 March 2013

**References**

- Balalaie S, Mahdidoust M, Eshaghi-Najafabadi R (2007) 2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate as an efficient coupling reagent for the amidation and phenylhydrazone of carboxylic acids at room temperature. *J Iran Chem Soc* 4:364–369
- Bunch L, Erichsen MN, Jensen AA (2009) Excitatory amino acid transporters as potential drug targets. *Expert Opin Ther Targets* 13:719–731
- Burkhard JA, Wagner B, Fischer H, Schuler F, Mueller K, Carreira EM (2010) Synthesis of azaspirocycles and their evaluation in drug discovery. *Angew Chem-Int Edit* 49:3524–3527
- Cook G, Barta N, Stille J (1992) Lewis acid-promoted 3-Aza-cope rearrangement of N-alkyl-N-allylenamines. *J Org Chem* 57:461–467
- Dunlop J, McIlvain HB, Carrick TA, Jow B, Lu Q, Kowal D, Lin S, Greenfield A, Grosanu C, Fan K, Petroski R, Williams J, Bastlund JF, Bundgaard C, Monrad O, Bekker-Jensen A, Nielsen CW, Frydenvang K, Jensen AA, Bunch L (2010) Characterization of novel aryl-ether, biaryl, and fluorene aspartic acid and diaminiopropionic acid analogs as potent inhibitors of the high-affinity glutamate transporter EAAT2. *Mol Pharmacol* 68:974–982
- Erichsen MN, Huynh THV, Abrahamsen B, Bastlund JF, Bundgaard C, Monrad O, Bekker-Jensen A, Nielsen CW, Frydenvang K, Jensen AA, Bunch L (2010) Structure–activity relationship study of first selective inhibitor of excitatory amino acid transporter subtype 1: 2-amino-4-(4-methoxyphenyl)-7-(naphthalen-1-yl)-5-oxo-5,6,7,8-tetrahydro-4 H -chromene-3-carbonitrile (UCPH-101). *J Med Chem* 53:7180–7191
- Fragasso G, Pallosi A, Puccetti P, Silipigni C, Rossodivita A, Pala M, Calori G, Alfieri O, Margonato A (2006) A randomized clinical trial of trimetazidine, a partial free fatty acid oxidation inhibitor, in patients with heart failure. *J Am Coll Cardiol* 48:992–998
- Gudipati M, Radziszewski J, Kaszynski P, Michl J (1993) Bicyclo[3.2.2]non-1-Ene - matrix-isolation and spectroscopic characterization of a moderately strained bridgehead olefin. *J Org Chem* 58:3668–3674
- Huynh THV, Shim I, Bohr H, Abrahamsen B, Nielsen B, Jensen AA, Bunch L (2012a) Structure–activity relationship study of selective excitatory amino acid transporter subtype 1 (EAAT1) inhibitor 2-amino-4-(4-methoxyphenyl)-7-(naphthalen-1-yl)-5-oxo-5,6,7,8-tetrahydro-4 H -chromene-3-carbonitrile (UCPH-101) and absolute configurational assignment using infrared and vibrational circular dichroism spectroscopy in combination with ab initio hartree–fock calculations. *J Med Chem* 55:5403–5412
- Huynh THV, Abrahamsen B, Madsen KK, Gonzalez-Franquesa A, Jensen AA, Bunch L (2012b) Design, synthesis and pharmacological characterization of coumarin-based fluorescent analogs of excitatory amino acid transporter subtype 1 selective inhibitors, UCPH-101 and UCPH-102. *Bioorg Med Chem* 20:6831–6839
- Jensen AA, Bräuner-Osborne H (2004) Pharmacological characterization of human excitatory amino acid transporters EAAT1, EAAT2 and EAAT3 in a fluorescence-based membrane potential assay. *Biochem Pharmacol* 67:2115–2127
- Jensen AA, Erichsen MN, Nielsen CW, Stensbøl TB, Kehrer J, Bunch L (2009) Discovery of the first selective inhibitor of excitatory amino acid transporter subtype 1. *J Med Chem* 52:912–915
- Lauriat TL, Richler E, McInnes LA (2007) A quantitative regional expression profile of EAAT2 known and novel splice variants reopens the question of aberrant EAAT2 splicing in disease. *Neurochem Int* 50:271–280
- Massie A, Vandesande F, Arckens L (2001) Expression of the high-affinity glutamate transporter EAAT4 in mammalian cerebral cortex. *Neuroreport* 12:393–397
- Millan MJ, Cussac D, Milligan G, Carr C, Audinot V, Gobert A, Lejeune F, Rivet JM, Brocco M, Duqueyroux D, Nicolas JP, Boutin JA, Newman-Tancredi A (2001) Antiparkinsonian agent piribedil displays antagonist properties at native, rat, and cloned, human  $\alpha$ (2)-adrenoceptors: cellular and functional characterization. *J Pharmacol Exp Ther* 297:876–887
- Nieouillon A, Canolle B, Masmejean F, Guillet B, Pisano P, Lortet S (2006) The neuronal excitatory amino acid transporter EAAC1/EAAT3: does it represent a major actor at the brain excitatory synapse? *J Neurochem* 98:1007–1018
- Nishiyama M, Yamamoto T, Koie Y (1998) Synthesis of N-arylpiperazines from aryl halides and piperazine under a palladium tri-tert-butylphosphine catalyst. *Tetrahedron Lett* 39:617–620



- Sagot E, Jensen AA, Pickering DS, Pu X, Umberti M, Stensbøl TB, Nielsen B, Assaf Z, Aboab B, Bolte J, Gefflaut T, Bunch L (2008) Chemo-enzymatic synthesis of (2S,4R)-2-amino-4-(3-(2,2-diphenylethylamino)-3-oxopropyl)pentanedioic acid: a novel selective inhibitor of human excitatory amino acid transporter subtype 2. *J Med Chem* 51:4085–4092
- Weisberg E, Manley PW, Cowan-Jacob SW, Hochhaus A, Griffin JD (2007) Second generation inhibitors of BCR-ABL for the treatment of imatinib-resistant chronic myeloid leukaemia. *Nat Rev Cancer* 7:345–356

doi:10.1186/2193-1801-2-112

**Cite this article as:** Huynh et al.: Structure-activity-relationship study of *N*-acyl-*N*-phenylpiperazines as potential inhibitors of the Excitatory Amino Acid Transporters (EAATs): improving the potency of a micromolar screening Hit is not truism. *SpringerPlus* 2013 2:112.

**Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:**

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

---

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)

---