

RESEARCH

Open Access



A range division and contraction approach for nonconvex quadratic program with quadratic constraints

Chunshan Xue^{1*}, Hongwei Jiao², Jingben Yin² and Yongqiang Chen³

*Correspondence:
xuechunshan2015@126.com

¹ School of Mathematics
and Statistics, Zhoukou
Normal University,
Zhoukou 466001, China
Full list of author information
is available at the end of the
article

Abstract

This paper presents a novel range division and contraction approach for globally solving nonconvex quadratic program with quadratic constraints. By constructing new underestimating linear relaxation functions, we can transform the initial nonconvex quadratic program problem into a linear program relaxation problem. By employing a branch and bound scheme with a range contraction approach, we describe a novel global optimization algorithm for effectively solving nonconvex quadratic program with quadratic constraints. Finally, the global convergence of the proposed algorithm is proved and numerical experimental results demonstrate the effectiveness of the proposed approach.

Keywords: Nonconvex quadratic program with quadratic constraints, Global optimization, Underestimating linear relaxation, Range contraction approach, Branch and bound scheme

Mathematics Subject Classification: 90C20, 90C26, 65K05

Background

The mathematical modeling of nonconvex quadratic program with quadratic constraints can be formulated as follows:

$$(NQPQC) : \begin{cases} \min F_0(x) = x^T Q_0 x + d_0^T x \\ \text{s.t. } F_i(x) = x^T Q_i x + d_i^T x \leq b_i, \quad i = 1, \dots, m, \\ x \in X^0 = \{x \in R^n : l^0 \leq x \leq u^0\}, \end{cases}$$

where $Q_i, i = 0, 1, \dots, m$, are all symmetric $n \times n$ matrices, $d_0, d_i \in R^n, b_i \in R, i = 1, \dots, m; l^0 = (l_1^0, \dots, l_n^0)^T, u^0 = (u_1^0, \dots, u_n^0)^T$.

Nonconvex quadratic program with quadratic constraints is worthy of study. On the one hand, this is because the kind of problems have many applications in practical problems, such as, heat exchanger engineering design, financial optimization, image processing, management science, etc (see Floudas and Visweswaran 1995; Shen 2007; Horst and Tuy 1996; Jiao and Liu in press; Bajirov and Rubinov 2001; Konno and Wiyayanayake 2001; Sherali and Smith 1997). On the other hand, it is since a large number of nonlinear

programming problems can be converted into this mathematical modeling, and the solutions of many nonlinear optimization problems can be approximated or obtained by solving a sequence of nonconvex quadratic programs with quadratic constraints. Moreover, from research point of view, since the nonconvex quadratic programs with quadratic constraints possess multiple local minimum points which are not globally minimum point, they exist significant computational and theoretical challenges. Therefore, it is necessary to put forward a new global optimization approach for solving the nonconvex quadratic program with quadratic constraints.

In past several decades, various algorithms have been developed for solving the nonconvex quadratic program with quadratic constraints and its special form, which are given as follows. Based on Newton method, branching rule and cutting plane, Vandenbussche and Nemhauser presented a branch-and-cut method for nonconvex quadratic programs with box constraints (Vandenbussche and Nemhauser 2005). Used different d.c. decomposition method to construct the relaxation of quadratic objective function, and used the “optimal level solution” parametrical approach to solve relaxation problem, Cambini and Sodini (2005) proposed a decomposition method for solving nonconvex quadratic programs over a compact polyhedral feasible region. By decomposing a large-scale quadratic programming into a serial of small-scale ones and approximating the solution of the large-scale quadratic programming via the solutions of these small-scale ones, Li and Zhang (2006) presented a decomposition algorithm for solving large-scale quadratic programming problems. Through decomposing quadratic objective function into a separable equivalent function, then by constructing linear under-estimator of the corresponding objective function, Shen et al. (2008) presented a decomposition and linearization method for globally solving quadratic programs with linear constraints. Vavasis (1992) presented an approximation algorithm for indefinite quadratic programming, and concluded that such an approximation solution can be found in polynomial time. Based on D.C. decomposition, Cholesky factorization and convex relaxation, Yajima and Fujie (1998) proposed a decomposition-and-relaxation algorithm for the general quadratic problems with box constraints. Ye (1992) proposed an affine-scaling algorithm to solve nonconvex indefinite or negative definite quadratic programs problems. By utilizing cutting plane technique, Gao and Deng (2008) presented a branch and bound method mixed with cutting plane technique for solving concave quadratic programming problems. Using lagrangian underestimates and interval Newton method, Voorhis (2002) proposed a global optimization algorithm for quadratic programs. Based on simplicial branch-and-bound scheme, Raber (1998) presented a simplicial branch-and-bound method for solving nonconvex quadratic programs. Based on parametric linear relaxation and new linearizing technique, Jiao et al. (2015), Jiao and Chen (2013) proposed two branch and bound algorithms for globally solving nonconvex quadratic programs. Using duality bounds and branch-and-bound scheme, Thoai (2000) presented a duality bound method for the general quadratic programming problem with quadratic constraints. Based on linear relaxation method and branch-and-bound framework with rectangle reducing technique, Shen and Liu (2008), Gao et al. (2005a, b), Jiao et al. (2014) proposed four branch-and-reduce algorithms for solving nonconvex quadratic programs, respectively. Based on branch-and-bound scheme, An and Tao (1997), An and Tao (1998) presented two D.C. algorithms for solving nonconvex quadratic programs.

Apart from the above reviewed quadratic programs algorithms, many algorithms (see Shen and Jiao 2006; Wang and Liang 2005; Wang et al. 2004; Shen 2005; Shen and Li 2013; Shen and Bai 2013; Shen et al. 2009) for solving geometric programming can be also used to solve the nonconvex quadratic program with quadratic constraints proposed in this paper. In addition, some recent artificial intelligent optimization algorithms (Zhang et al. 2013, 2014a, b, 2016) are developed, which are also used to obtain local optimal solution of the nonconvex quadratic program with quadratic constraints.

The above most deterministic algorithms for solving nonconvex quadratic programs are based on relaxation technique and branch-and-bound scheme, since the exhaustiveness of branching rule leads to a significant increase in the computational burden for solving nonconvex quadratic programs, until today, there is short of more effective algorithm for solving such problems, so it is necessary to establish a good algorithm for the NQPQC. Therefore, the main motivation for the paper is to construct a novel linearized technique and a range contraction approach, and based on these techniques develop an effective algorithm for solving the NQPQC.

In this paper, a novel range division and contraction approach will be proposed for globally solving the nonconvex quadratic program with quadratic constraints. The main features of the proposed algorithm in this paper are as follows. Firstly, a novel linearized technique is constructed for systematically converting the NQPQC into a sequence of linear program problems, and the solutions of the converted linear program problems are used to approximate the solution of the NQPQC, which can be computed by using simplex approach. Secondly, in order to accelerate the running speed of our approach, a range contraction method is constructed and employed in our algorithm. Next, use branch-and-bound framework with a range contraction approach, a new global optimization algorithm is established. Finally, the global convergence of the proposed approach is proved and numerical experimental results demonstrate the computational efficiency of our algorithm.

The remaining contents of the paper are stated as follows. Second section describes a new linearized method and the linear program relaxation problem of the initial NQPQC is derived. In third section, a new range division and contraction approach is established and its global convergence is proved. Fourth section describes numerical results for some examples, which be computed by using the proposed algorithm. Finally some conclusions are drawn.

New linearized approach

In this section, a new linearized approach is constructed for establishing the linear program relaxation problem of the NQPQC. Without loss of generality, let $X^k = \{x \in R^n \mid l^k = (l_1^k, \dots, l_n^k)^T \leq x \leq u^k = (u_1^k, \dots, u_n^k)^T\} \subseteq X^0$, and let λ_i be the minimum eigenvalue of Q_i . Set

$$\theta_i = \begin{cases} 0, & \text{if } \lambda_i \geq 0, \\ |\lambda_i| + \rho, & \text{if } \lambda_i < 0, \end{cases} \text{ where } \rho \text{ is an arbitrary positive real number.}$$

In the following, for any $i \in \{0, 1, \dots, m\}$, for any $x \in X^k$, define

$$F_i^L(x) = \left[d_i + Q_i^T (l^k + u^k) + \theta_i (u^k - l^k) \right]^T x + 2\theta_i (u^k)^T l^k - \theta_i (u^k)^T (u^k) - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right).$$

Theorem 1 For any $x \in X^k$, consider the functions $F_i(x)$ and $F_i^L(x)$, we have:

- (i) $F_i^L(x) \leq F_i(x), i = 0, 1, \dots, m;$
- (ii) For each $i = 0, 1, \dots, m, \|F_i(x) - F_i^L(x)\| \rightarrow 0$ as $\|u^k - l^k\| \rightarrow 0$.

Proof (1) Consider the function x_j^2 over $[l_j^k, u_j^k]$, according to the mean value theorem, we can get

$$x_j^2 = 2\xi_j^k (x_j - u_j^k) + (u_j^k)^2 \leq 2l_j^k (x_j - u_j^k) + (u_j^k)^2 = 2l_j^k x_j - 2u_j^k l_j^k + (u_j^k)^2,$$

where $\xi_j^k = \alpha l_j^k + (1 - \alpha)u_j^k, \alpha \in [0, 1], j = 1, \dots, n$.

By the expressions of the functions $F_i(x)$ and $F_i^L(x)$, we can get that

$$\begin{aligned} F_i(x) &= x^T Q_i x + d_i^T x \\ &= \left(x - \frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(x - \frac{l^k + u^k}{2} \right) \\ &\quad + d_i^T x - \theta_i \sum_{j=1}^n x_j^2 + (l^k + u^k)^T (Q_i + \theta_i I) x - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right) \\ &\geq d_i^T x - \theta_i \sum_{j=1}^n x_j^2 + (l^k + u^k)^T (Q_i + \theta_i I) x - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right) \\ &\geq d_i^T x + \theta_i \sum_{j=1}^n \left[-2l_j^k x_j + 2u_j^k l_j^k - (u_j^k)^2 \right] + (l^k + u^k)^T (Q_i + \theta_i I) x \\ &\quad - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right) \\ &= \left[d_i + (Q_i + \theta_i I)^T (l^k + u^k) - 2\theta_i l^k \right]^T x + 2\theta_i (u^k)^T l^k - \theta_i (u^k)^T (u^k) \\ &\quad - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right) \\ &= \left[d_i + Q_i^T (l^k + u^k) + \theta_i (u^k - l^k) \right]^T x + 2\theta_i (u^k)^T l^k - \theta_i (u^k)^T (u^k) \\ &\quad - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right) \\ &= F_i^L(x). \end{aligned}$$

Therefore, we have

$$F_i^L(x) \leq F_i(x), \quad i = 0, 1, \dots, m.$$

(2) Consider $F_i(x)$ and $F_i^L(x)$, let $\Delta_i = \|F_i(x) - F_i^L(x)\|$, we can follow that

$$\begin{aligned} \Delta_i &= \left\| x^T Q_i x + d_i^T x - \left\{ \left[d_i + Q_i^T (l^k + u^k) + \theta_i (u^k - l^k) \right]^T x + 2\theta_i (u^k)^T l^k \right. \right. \\ &\quad \left. \left. - \theta_i (u^k)^T (u^k) - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right) \right\} \right\| \\ &= \left\| \left[\left(x - \frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(x - \frac{l^k + u^k}{2} \right) + d_i^T x - \theta_i \sum_{j=1}^n x_j^2 \right. \right. \\ &\quad \left. \left. + (l^k + u^k)^T (Q_i + \theta_i I) x - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right) \right] \right. \\ &\quad \left. - \left\{ \left[d_i + (Q_i + \theta_i I)^T (l^k + u^k) - 2\theta_i l^k \right]^T x + 2\theta_i (u^k)^T l^k \right. \right. \\ &\quad \left. \left. - \theta_i (u^k)^T (u^k) - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right) \right\} \right\| \\ &= \left\| \left(x - \frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(x - \frac{l^k + u^k}{2} \right) \right. \\ &\quad \left. + \theta_i \sum_{j=1}^n \left[2l_j^k x_j - 2u_j^k l_j^k + (u_j^k)^2 - x_j^2 \right] \right\| \\ &= \left\| \left(x - \frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(x - \frac{l^k + u^k}{2} \right) \right. \\ &\quad \left. + \theta_i \sum_{j=1}^n (u_j^k - x_j) (x_j - l_j^k + u_j^k - l_j^k) \right\| \\ &\leq \left\| \left(x - \frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(x - \frac{l^k + u^k}{2} \right) \right\| \\ &\quad + \theta_i \left\| \sum_{j=1}^n (u_j^k - x_j) (x_j - l_j^k + u_j^k - l_j^k) \right\| \\ &\leq \left\| \left(x - \frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(x - \frac{l^k + u^k}{2} \right) \right\| + \theta_i \left\| \sum_{j=1}^n (u_j^k - x_j) (x_j - l_j^k) \right\| \\ &\quad + \theta_i \left\| \sum_{j=1}^n (u_j^k - x_j) (u_j^k - l_j^k) \right\| \\ &\leq \left\| \left(x - \frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(x - \frac{l^k + u^k}{2} \right) \right\| + 2\theta_i \left\| (u^k - l^k)^T (u^k - l^k) \right\| \\ &= \frac{1}{4} \|Q_i + \theta_i I\| \|u^k - l^k\|^2 + 2\theta_i \|u^k - l^k\|^2 \end{aligned}$$

Since

$$\frac{1}{4} \|Q_i + \theta_i I\| \|u^k - l^k\|^2 + 2\theta_i \|u^k - l^k\|^2 \rightarrow 0 \text{ as } \|u^k - l^k\| \rightarrow 0,$$

therefore, we get

$$\Delta_i = \|F_i(x) - F_i^L(x)\| \rightarrow 0 \text{ as } \|u^k - l^k\| \rightarrow 0.$$

The conclusion is drawn. \square

From the above Theorem 1, we can establish the linear program relaxation problem (LPRP) of the NQPQC in X^k as follows.

$$\text{LPRP}(X^k) : \begin{cases} \min F_0^L(x) \\ \text{s.t. } F_i^L(x) \leq b_i, \quad i = 1, \dots, m, \\ x \in X^k = \{x : l^k \leq x \leq u^k\}. \end{cases}$$

where

$$\begin{aligned} F_i^L(x) = & \left[d_i + Q_i^T (l^k + u^k) + \theta_i (u^k - l^k) \right]^T x + 2\theta_i (u^k)^T l^k - \theta_i (u^k)^T (u^k) \\ & - \left(\frac{l^k + u^k}{2} \right)^T (Q_i + \theta_i I) \left(\frac{l^k + u^k}{2} \right). \end{aligned}$$

Obviously, by the construction method of linear program relaxation problem (LPRP), we get that the feasible region of the LPRP contain all feasible points of the NQPQC in X^k , and the optimal value of the LPRP is not more than that of the NQPQC in X^k .

Range contraction approach

To accelerate running speed of our algorithm, a range contraction approach is formulated as the following Theorem 2. The proposed range contraction approach aims at contracting the investigated rectangle X without pruning any global optimum point of the initial NQPQC.

Without loss of generality, for any $x \in X = (X_j)_{n \times 1} \subseteq X^0$ with $X_j = [l_j, u_j]$ ($j = 1, \dots, n$), we can rewrite the function $F_i^L(x)$ as follows:

$$F_i^L(x) = \sum_{j=1}^n c_{ij} x_j + \delta_i, \quad i = 0, 1, \dots, m.$$

Let \overline{UB} be a known currently upper bound of the global optimum value of the NQPQC(X^0), and set

$$\begin{aligned}\widehat{LB}_i &= \sum_{j=1}^n \min\{c_{ij}l_j, c_{ij}u_j\} + \delta_i, \quad i = 0, 1, \dots, m, \\ \overline{X}_j &= \begin{cases} X_j, & j \in \{1, \dots, p-1, p+1, \dots, n\}, \\ \left[\frac{\overline{UB} - \widehat{LB}_0 + \min\{c_{0p}l_p, c_{0p}u_p\}}{c_{0p}}, u_p \right] \cap X_p, & j = p; \end{cases} \\ \underline{X}_j &= \begin{cases} X_j, & j \in \{1, \dots, p-1, p+1, \dots, n\}, \\ \left[l_p, \frac{\overline{UB} - \widehat{LB}_0 + \min\{c_{0p}l_p, c_{0p}u_p\}}{c_{0p}} \right] \cap X_p, & j = p; \end{cases} \\ \tilde{X}_j &= \begin{cases} X_j, & j \in \{1, \dots, p-1, p+1, \dots, n\}, \\ \left[\frac{b_i - \widehat{LB}_i + \min\{c_{ip}l_p, c_{ip}u_p\}}{c_{ip}}, u_p \right] \cap X_p, & j = p; \end{cases} \\ \hat{X}_j &= \begin{cases} X_j, & j \in \{1, \dots, p-1, p+1, \dots, n\}, \\ \left[l_p, \frac{b_i - \widehat{LB}_i + \min\{c_{ip}l_p, c_{ip}u_p\}}{c_{ip}} \right] \cap X_p, & j = p. \end{cases}\end{aligned}$$

Theorem 2 For any sub-rectangle $X \subseteq X^0$, the following conclusions hold:

- If $\widehat{LB}_0 > \overline{UB}$ or $\widehat{LB}_i > b_i$ for some $i \in \{1, \dots, m\}$, then there contains no global optimal solution of the NQPQC(X^0) over X .
- If $\widehat{LB}_0 \leq \overline{UB}$, then, for any $p \in \{1, 2, \dots, n\}$, if $c_{0p} > 0$, then there does not contain global optimal solution of the NQPQC(X^0) over $\overline{X} = (\overline{X}_j)_{n \times 1}$; if $c_{0p} < 0$, there does not contain global optimal solution of the NQPQC(X^0) over $\underline{X} = (\underline{X}_j)_{n \times 1}$.
- If $\widehat{LB}_i \leq b_i$ for each $i \in \{1, \dots, m\}$, then, for any $p \in \{1, 2, \dots, n\}$, if $c_{ip} > 0$, then there does not contain global optimal solution of the NQPQC(X^0) over $\tilde{X} = (\tilde{X}_j)_{n \times 1}$; if $c_{ip} < 0$, there does not contain global optimal solution of the NQPQC(X^0) over $\hat{X} = (\hat{X}_j)_{n \times 1}$.

Proof The proof of the Theorem can be similarly given by the Theorem 3 in Jiao et al. (2014), therefore, here is omitted. \square

From the above Theorem 2, a new range contraction approach is presented as follows:

Range contraction approach

Without loss of generality, assume that $X = (X_j)_{n \times 1}$ with $X_j = [l_j, u_j]$ ($j = 1, \dots, n$) be any sub-rectangle of X^0 , and for each $i = 0, 1, \dots, M$, compute \widehat{LB}_i .

If there exist some $i \in \{1, \dots, M\}$, such that $\widehat{LB}_i > b_i$, then delete the whole rectangle X ;

Otherwise, for each $i \in \{1, \dots, M\}$, $p \in \{1, \dots, n\}$, compute $\frac{b_i - \widehat{LB}_i + \min\{c_{ip}l_p, c_{ip}u_p\}}{c_{ip}}$. If $c_{ip} > 0$ and $\frac{b_i - \widehat{LB}_i + \min\{c_{ip}l_p, c_{ip}u_p\}}{c_{ip}} < u_p$, then let $u_p = \frac{b_i - \widehat{LB}_i + \min\{c_{ip}l_p, c_{ip}u_p\}}{c_{ip}}$; else if $c_{ip} < 0$ and $\frac{b_i - \widehat{LB}_i + \min\{c_{ip}l_p, c_{ip}u_p\}}{c_{ip}} > l_p$, then let $l_p = \frac{b_i - \widehat{LB}_i + \min\{c_{ip}l_p, c_{ip}u_p\}}{c_{ip}}$.

If $\widehat{LB}_0 > \overline{UB}$, then delete the whole rectangle X ;

Otherwise, for each $p \in \{1, \dots, n\}$, compute $\frac{\overline{UB} - \widehat{LB}_0 + \min\{c_{0p}l_p, c_{0p}u_p\}}{c_{0p}}$. If $c_{0p} > 0$ and $\frac{\overline{UB} - \widehat{LB}_0 + \min\{c_{0p}l_p, c_{0p}u_p\}}{c_{0p}} < u_p$, then let $u_p = \frac{\overline{UB} - \widehat{LB}_0 + \min\{c_{0p}l_p, c_{0p}u_p\}}{c_{0p}}$; else if $c_{0p} < 0$ and $\frac{\overline{UB} - \widehat{LB}_0 + \min\{c_{0p}l_p, c_{0p}u_p\}}{c_{0p}} > l_p$, then let $l_p = \frac{\overline{UB} - \widehat{LB}_0 + \min\{c_{0p}l_p, c_{0p}u_p\}}{c_{0p}}$.

From Theorem 2, by utilizing the above range contraction approach to compress the investigated rectangle region, or delete a part of the investigated rectangle region, in which there does not contain the global optimum solution of the NQPQC. Therefore, we can improve the computational speed or computational efficiency of the algorithm.

Range division and contraction algorithm

In this section, a new range division and contraction algorithm is presented for globally solving the initial NQPQC. In the algorithm, one of the most important operation is the choice of a suitable range division method. Here, we choose an w -division approach, which is sufficient to ensure the global convergence of the proposed algorithm, this is because that the selected range division method drives all interval to zero for all variables. This range division method is described as follows.

For any investigated rectangle $X' = [\underline{x}', \bar{x}'] \subseteq X^0$. Denote $q \in \arg \max \{\bar{x}'_i - \underline{x}'_i : i = 1, 2, \dots, n\}$, and by dividing the interval $[\underline{x}'_q, \bar{x}'_q]$ into two new subintervals $[\underline{x}'_q, \underline{x}'_q + \omega(\bar{x}'_q - \underline{x}'_q)]$ and $[\underline{x}'_q + \omega(\bar{x}'_q - \underline{x}'_q), \bar{x}'_q]$, where $\omega \in (0, 1)$. At the same time, all other interval remain unchanged. Consequently, the investigated rectangle X' is partitioned into two new sub-rectangles X'_1 and X'_2 .

Without loss of generality, we denote $LB(X^k)$ and $x^k = x(X^k)$ as the optimum value and optimal solution of the linear program relaxation problem over the sub-rectangle X^k , respectively. Combining the former linear program relaxation problem, range division method and range contraction approach together, a new global optimization method is constructed for effectively solving the NQPQC, the main steps of the proposed algorithm are described as follows.

Range division and contraction algorithm

Step 0. (Initializing) Set the initial iteration number $k = 0$, the initial active node set $\Delta_0 = \{X^0\}$, the given termination error $\epsilon > 0$, the initial upper bound $UB_0 = +\infty$ and the initial feasible point set $\Lambda = \emptyset$, respectively.

Step 1. (Judgement) Compute the LPRP(X^0), denote its optimum solution $x^0 = x(X^0)$ and optimum value $LB(X^0)$, respectively. If x^0 is a feasible point of the NQPQC(X^0), let $UB_0 = F_0(x^0)$ and $\Lambda = \Lambda \cup \{x^0\}$. Let $LB_0 = LB(X^0)$, if $UB_0 - LB_0 \leq \epsilon$, then x^0 is a global optimum solution of the NQPQC(X^0). Otherwise, proceed to Step 2.

Step 2. (Division) Use the proposed range division method to divide X^k into two new sub-rectangles, and denote the set of new partitioned sub-rectangle as \bar{X}^k .

Step 3. (Contraction) For each investigated sub-rectangle $X \in \bar{X}^k$, use the presented range contraction approach to compress its range, and still denote the remaining rectangle part and the remaining partitioning set by X and \bar{X}^k , respectively.

Step 4. (Bounding) If \bar{X}^k is not empty, for each $X \in \bar{X}^k$, compute the LPRP(X), and denote its optimum value and optimum solution by $LB(X)$ and $x(X)$, respectively. If $LB(X) > UB_k$, let $\bar{X}^k := \bar{X}^k \setminus X$; else if the midpoint x^{mid} of X is a feasible solution of the NQPQC(X^0), then let $\Lambda := \Lambda \cup \{x^{mid}\}$, and if $x(X)$ is a feasible solution of the NQPQC(X^0), then let $\Lambda := \Lambda \cup \{x(X)\}$.

If Λ is not empty, denote the new upper bound $UB_k := \min_{x \in \Lambda} F_0(x)$, and denote the best feasible point $x^* := \operatorname{argmin}_{x \in \Lambda} F_0(x)$.

Denote the new remaining partition set and the new lower bound by $\Theta_k := (\Theta_k \setminus X^k) \cup \bar{X}^k$ and $LB_k := \inf_{X \in \Theta_k} LB(X)$, respectively.

Step 5. (Judgement) Denote $\Theta_{k+1} = \Theta_k \setminus \{X : UB_k - LB(X) \leq \epsilon, X \in \Theta_k\}$. If $\Theta_{k+1} = \emptyset$, then we have: UB_k and x^* be the global optimum value and the global optimum solution of the initial NQPQC, respectively. Otherwise, select a new sub-rectangle X^{k+1} which satisfies $X^{k+1} = \operatorname{argmin}_{X \in \Theta_{k+1}} LB(X)$, let $k := k + 1$ and return to Step 2.

The above algorithm either terminates after finite iteration or generates an infinite iteration sequence, from the exhaustiveness of the used range division approach, we can follow that all intervals of all variables must shrink to a singleton, i.e., $\|u^k - l^k\| \rightarrow 0$. At the same time, the Theorem 1 guarantee that the linear program relaxation problem (LPRP) (X^k) will infinitely approaches the problem NQPQC(X^k) as $\|u^k - l^k\| \rightarrow 0$.

Theorem 3 *The above algorithm either finishes finitely at the global optimum point x^* of the initial NQPQC, or produces an infinite iteration sequence $\{x^k\}$, of which any accumulation point will be the global optimum point of the initial NQPQC.*

Proof If the above algorithm finishes finitely at some iteration k , obviously, when the algorithm ends, we can get $UB_k = v^* = LB_k$. Therefore, from step of the algorithm, we get that x^k is a global optimum point of the initial NQCQP.

If the above algorithm does not finish at finite step, then it must produce an infinite sub-rectangle iteration sequence $\{X^k\}$, from the exhaustiveness of division approach, we get that the sub-rectangle sequence $\{X^k\}$ must converge to a point. From the characteristics of our algorithm, we can get that $\{UB_k\}$ and $\{LB_k\}$ are nonincreasing and non-decreasing sequences, respectively. So that $\{UB_k - LB_k\}$ is a monotonic non-increasing sequence. By the conclusion of Theorem 1, we get that the sequence $\{UB_k - LB_k\}$ must be convergent to zero. From the structure of our algorithm, for each k , we have $LB_k \leq v^* \leq UB_k$. Therefore, we can get that $\lim_{k \rightarrow \infty} UB_k = \lim_{k \rightarrow \infty} LB_k = v^*$. From steps of our algorithm, we know that x^k is always a feasible point of the initial NQCQP, we can get $UB_k = G_0(x^k)$. By the continuity of constrained functions, we get that the limitation point x^* of $\{x^k\}$ is also a feasible point of the initial NQCQP, and we can get the global optimum value $v^* = G_0(x^*)$. Hence, the conclusion is proved. \square

Numerical experiments

In this section, in order to verify the performance of the algorithm proposed in this paper (our algorithm), several numerical examples in recent references are implemented on a Intel(R) Core(TM)2 Duo CPU microcomputer. Although these numerical examples

have a relatively small number of variables, they are also quite challenging. The proposed algorithm program is coded in C++ language, all linear program relaxation problems are solved by using simplex approach, and the convergence errors are all set as $\epsilon = 10^{-6}$ in our experiments. For all examples, the numerical results of optimal solutions, optimal values and number of iterations obtained by our algorithm and other approaches (Jiao and Chen 2013; Thoai 2000; Shen and Liu 2008; Gao et al. 2005b; Jiao et al. 2014; Shen and Jiao 2006; Wang and Liang 2005; Wang et al. 2004; Shen 2005; Shen and Li 2013) are illustrated in Table 1. The numerical experimental results show that our algorithm can globally solve the NQCQP problem. In Table 1, the notation “Iter.” represents “number of iterations”.

Table 1 Numerical comparisons with the known approaches for Examples 1–8

Example	References	Optimum value	Optimum solution	Iter.
1	This paper	−16.000000000	(5.000000000, 1.000000000)	5
	Thoai (2000)	−16.0	(5.0, 1.0)	10
	Jiao and Chen (2013)	−16.000000000	(5.000000000, 1.000000000)	5
2	This paper	6.777777779	(2.000000000, 1.666666667)	14
	Shen and Liu (2008)	6.777782016	(2.000000000, 1.666666667)	40
	Jiao et al. (2014)	6.77778233	(2.0, 1.666666667)	32
	Jiao et al. (2014)	6.77778517	(2.0, 1.666666667)	33
	Wang and Liang (2005)	6.7780	(2.00003, 1.66665)	44
3	This paper	0.500000394	(0.500000000, 0.500000000)	28
	Shen and Liu (2008)	0.500004627	(0.5 0.5)	34
	Wang and Liang (2005)	0.5	(0.5, 0.5)	91
	Wang et al. (2004)	0.5	(0.5, 0.5)	96
	Jiao and Chen (2013)	0.500000442	(0.500000000, 0.500000000)	37
4	This paper	118.383672094	(2.555940349, 3.129963478)	45
	Gao et al. (2005b)	118.383756475	(2.5557793695, 3.13016463929)	210
	Jiao et al. (2014)	118.383672231	(2.555800904, 3.130134268)	61
	Jiao and Chen (2013)	118.383671904	(2.555745855, 3.130201688)	59
5	This paper	−1.162881826	(1.500000000, 1.500000000)	31
	Shen (2005)	−1.16288	(1.5, 1.5)	84
	Jiao and Chen (2013)	−1.162882693	(1.500000000, 1.500000000)	24
6	This paper	1.177125219	(1.177124344, 2.177124344)	12
	Shen and Jiao (2006)	1.177124327	(1.177124327, 2.177124353)	434
	Jiao and Chen (2013)	1.177125051	(1.177124344, 2.177124344)	22
7	This paper	−0.999999102	(2.000000000, 1.000000000)	20
	Shen and Jiao (2006)	−1.0	(2.000000, 1.000000)	24
	Jiao and Chen (2013)	−0.999999410	(2.000000000, 1.000000000)	21
8	This paper	−11.36363579	(1.0, 0.181818133, 0.983332175)	352
	Shen and Li (2013)	−10.35	(0.998712, 0.196213, 0.979216)	1648
	Jiao and Chen (2013)	−11.363636364	(1.0, 0.181818470, 0.983332113)	420

Example 1 (Jiao and Chen 2013; Thoai 2000).

$$\begin{cases} \min & -x_1^2 + x_1x_2 + x_2^2 + x_1 - 2x_2 \\ \text{s.t.} & x_1 + x_2 \leq 6, \\ & -2x_1^2 + x_2^2 + 2x_1 + x_2 \leq -4, \\ & 1 \leq x_1 \leq 6, \quad 1 \leq x_2 \leq 6. \end{cases}$$

For Example 1, from numerical results in Table 1, compared with the algorithms in Jiao and Chen (2013), Thoai (2000), use the same logic of the algorithm, our algorithm can obtain the same optimal solution (5.0, 1.0), but our algorithm spends less number of iterations.

Example 2 (Shen and Liu 2008; Jiao et al. 2014; Wang and Liang 2005).

$$\begin{cases} \min & x_1^2 + x_2^2 \\ \text{s.t.} & 0.3x_1x_2 \geq 1, \\ & 2 \leq x_1 \leq 5, \quad 1 \leq x_2 \leq 3. \end{cases}$$

For Example 2, by Table 1, compared with the algorithms in Shen and Liu (2008), Jiao et al. (2014), use the same logic of the algorithm, our algorithm can obtain the same optimal solution with less number of iterations; and compared with the algorithms in Wang and Liang (2005), our algorithm can obtain the better optimal solution with less number of iterations.

Example 3 (Jiao and Chen 2013; Shen and Liu 2008; Wang and Liang 2005; Wang et al. 2004).

$$\begin{cases} \min & x_1 \\ \text{s.t.} & 4x_2 - 4x_1^2 \leq 1, \\ & -x_1 - x_2 \leq -1, \\ & 0.01 \leq x_1 \leq 15, \quad 0.01 \leq x_2 \leq 15. \end{cases}$$

From the numerical results of Example 3 in Table 1, compared with the algorithms in Jiao and Chen (2013), Shen and Liu (2008), Wang and Liang (2005), Wang et al. (2004), our algorithm can obtain the same optimal solution (0.5, 0.5) as Jiao and Chen (2013), Shen and Liu (2008), Wang and Liang (2005), Wang et al. (2004), but our algorithm spends less number of iterations.

Example 4 (Jiao and Chen 2013; Gao et al. 2005b; Jiao et al. 2014).

$$\begin{cases} \min & 6x_1^2 + 4x_2^2 + 5x_1x_2 \\ \text{s.t.} & -6x_1x_2 \leq -48, \\ & 0 \leq x_1 \leq 10, \quad 0 \leq x_2 \leq 10. \end{cases}$$

For Example 4, by the numerical results in Table 1, compared with the algorithms in Jiao and Chen (2013), Gao et al. (2005b), Jiao et al. (2014), our algorithm can obtain better or at least as good as optimal solution and optimal value with less number of iterations.

Example 5 (Jiao and Chen 2013; Shen 2005).

$$\begin{cases} \min & -x_1 + x_1 x_2^{0.5} - x_2 \\ \text{s.t.} & -6x_1 + 8x_2 \leq 3, \\ & 3x_1 - x_2 \leq 3, \\ & 1 \leq x_1 \leq 1.5, 1 \leq x_2 \leq 1.5. \end{cases}$$

For Example 5, by the numerical results of Table 1, compared with the algorithms in Jiao and Chen (2013), Shen (2005), our algorithm can obtain the same optimal solution (1.5, 1.5), but our algorithm spend less number of iterations.

Example 6 (Jiao and Chen 2013; Shen and Jiao 2006).

$$\begin{cases} \min & x_1 \\ \text{s.t.} & \frac{1}{4}x_1 + \frac{1}{2}x_2 - \frac{1}{16}x_1^2 - \frac{1}{16}x_2^2 \leq 1, \\ & \frac{1}{14}x_1^2 + \frac{1}{14}x_2^2 - \frac{3}{7}x_1 - \frac{3}{7}x_2 \leq -1, \\ & 1 \leq x_1 \leq 5.5, 1 \leq x_2 \leq 5.5. \end{cases}$$

For Example 6, from the numerical results in Table 1, our algorithm obtains the optimal solution (1.177124344, 2.177124344) after 24 iterations, but the algorithm in Shen and Jiao (2006) obtains the optimal solution (1.177124327, 2.177124353) after 24 iteration, obviously our algorithm has higher computational efficiency than the algorithms in Shen and Jiao (2006); compared with the algorithm in Jiao and Chen (2013), our algorithm can obtain the same optimal solution (1.177124344, 2.177124344), but our algorithm spend less number of iterations.

Example 7 (Jiao and Chen 2013; Shen and Jiao 2006).

$$\begin{cases} \min & x_1 x_2 - 2x_1 + x_2 + 1 \\ \text{s.t.} & 8x_2^2 - 6x_1 - 16x_2 \leq -11, \\ & -x_2^2 + 3x_1 + 2x_2 \leq 7, \\ & 1 \leq x_1 \leq 2.5, \\ & 1 \leq x_2 \leq 2.225. \end{cases}$$

For Example 7, from the numerical results in Table 1, compared with the algorithms in Jiao and Chen (2013), Shen and Jiao (2006), our algorithm can obtain the same optimal solution (2.0, 1.0), but our algorithm spend less number of iterations, obviously our

algorithm has higher computational efficiency than the algorithms in Jiao and Chen (2013), Shen and Jiao (2006).

Example 8 (Jiao and Chen 2013; Shen and Li 2013).

$$\begin{cases} \min & -4x_2 + (x_1 - 1)^2 + x_2^2 - 10x_3^2 \\ \text{s.t.} & x_1^2 + x_2^2 + x_3^2 \leq 2, \\ & (x_1 - 2)^2 + x_2^2 + x_3^2 \leq 2, \\ & 2 - \sqrt{2} \leq x_1 \leq \sqrt{2}, \\ & 0 \leq x_2, x_3 \leq \sqrt{2}. \end{cases}$$

For Example 8, from the numerical results in Table 1, our algorithm obtains the optimal solution (1.0, 0.181818133, 0.983332175) and the optimal value -11.363635790 after 352 iterations, but the algorithm in Jiao and Chen (2013) obtains the optimal solution (1.0, 0.181818470, 0.983332113) and optimal value -11.363636364 after 420 iteration, and the algorithm in Shen and Li (2013) obtains the optimal solution (0.998712, 0.196213, 0.979216) and optimal value -10.35 after 1648 iteration, obviously our algorithm not only has higher efficiency but also get the better optimal solution and optimal value.

In all, from numerical results for Examples 1–8, compared with Jiao and Chen (2013), Thoai (2000), Shen and Liu (2008), Gao et al. (2005b), Jiao et al. (2014), Shen and Jiao (2006), Wang and Liang (2005), Wang et al. (2004), Shen (2005), Shen and Li (2013), the presented algorithm in this paper can globally solve nonconvex quadratic program with quadratic constraints.

Conclusion

In this article, a new range division and contraction algorithm is proposed for globally solving nonconvex quadratic program with quadratic constraints (NQPQC). The linear program relaxation problem of the initial NQPQC is constructed by utilizing new linearizing method, which is derived by underestimating all quadratic objective function and constraint functions with linear relaxation functions. By applying the current upper bound and linear program relaxation problem of the NQPQC, a range contraction technique is introduced for improving the computational speed of our algorithm. By successive partition of the initial rectangle region, and by subsequently solving a sequence of linear program relaxation problems, the proposed algorithm converges to the global optimum point of the original NQPQC. Finally, Numerical computational results demonstrate the effectiveness and robustness of our algorithm.

Authors' contributions

This work was carried out in collaboration with all authors. All authors have a good contribution to design the new range division and contraction algorithm, and to perform the numerical analysis of this research work. All authors read and approved the final manuscript.

Author details

¹ School of Mathematics and Statistics, Zhoukou Normal University, Zhoukou 466001, China. ² School of Mathematical Sciences, Henan Institute of Science and Technology, Xinxiang 453003, China. ³ College of Mathematics and Information Science, Henan Normal University, Xinxiang 453007, China.

Acknowledgements

This paper is supported by the National Natural Science Foundation of China under Grant (11171094), the Basic and Advanced Technology Research Project of Henan Province (152300410097,142300410464), the Science and Technology Key Project of Education Department of Henan Province (14A110024,16A110014), the Major Scientific Research Projects of Henan Institute of Science and Technology (2015ZD07), the National Natural Science Foundation of Henan Province (152300410097).

Competing interests

The authors declare that they have no competing interests.

Received: 9 February 2016 Accepted: 30 June 2016

Published online: 12 July 2016

References

- An LTH, Tao PD (1997) Solving a class of linearly constrained indefinite quadratic problems by DC algorithm. *J Glob Optim* 11:253–285
- An LTH, Tao PD (1998) A branch and bound method via DC optimization algorithms and ellipsoidal technique for box constrained nonconvex quadratic problems. *J Glob Optim* 13:171–206
- Bajirov AM, Rubinov AM (2001) Global optimization of marginal functions with applications to economic equilibrium. *J Glob Optim* 20:215–237
- Cambini R, Sordini C (2005) Decomposition methods for solving nonconvex quadratic programs via branch and bound. *J Glob Optim* 33:313–336
- Floudas CA, Visweswaran V (1995) Quadratic optimization. In: Horst R, Pardalos PM (eds) *Nonconvex optimization and its applications*, vol 2. Kluwer Academic Publishers, Dordrecht, pp 217–269
- Gao Y, Xue H, Shen P (2005a) A new rectangle branch-and-reduce approach for solving nonconvex quadratic programming problems. *Appl Math Comput* 168(2):1409–1418
- Gao Y, Shang Y, Zhang L (2005b) A branch and reduce approach for solving nonconvex quadratic programming problems with quadratic constraints. *OR Trans* 9(2):9–20
- Gao Y, Deng G (2008) A branch and bound method mixed with cutting plane technique for solving concave quadratic programming problems. *Chin J Eng Math* 25(4):589–596
- Horst R, Tuy H (1996) *Global optimization: deterministic approaches*. Springer, Berlin
- Jiao H, Chen Y (2013) A global optimization algorithm for generalized quadratic programming. *J Appl Math* 2013:1–9
- Jiao H, Liu S (in press) Range division and compression algorithm for quadratically constrained sum of quadratic ratios
- Jiao H, Chen Y, Cheng W (2014) A novel optimization method for nonconvex quadratically constrained quadratic programs. *Abstr Appl Anal* 2014:1–11
- Jiao H, Liu S, Lu N (2015) A parametric linear relaxation algorithm for globally solving nonconvex quadratic programming. *Appl Math Comput* 250:973–985
- Konno H, Wiyayanayake A (2001) Portfolio optimization problem under concave transaction costs and minimal transaction unit constraints. *Math Program* 89:233–250
- Li HM, Zhang KC (2006) A decomposition algorithm for solving large-scale quadratic programming problems. *Appl Math Comput* 173:394–403
- Raber U (1998) A simplicial branch-and-bound method for solving nonconvex all-quadratic programs. *J Glob Optim* 13:417–432
- Shen P (2005) Linearization method of global optimization for generalized geometric programming. *Appl Math Comput* 162:353–370
- Shen P (2007) *Global optimization method*. Science Press, Beijing
- Shen P, Bai X (2013) Global optimization for generalized geometric programming problems with discrete variables. *Optimization* 62:895–917
- Shen P, Jiao H (2006) A new rectangle branch-and-pruning approach for generalized geometric programming. *Appl Math Comput* 183:1027–1038
- Shen P, Li X (2013) Branch-reduction-bound algorithm for generalized geometric programming. *J Glob Optim* 56(3):1123–1142
- Shen P, Liu L (2008) A global optimization approach for quadratic programs with nonconvex quadratic constraints. *Chin J Eng Math* 25:923–926
- Shen P, Pei Y, Gu M (2008) A decomposition and linearization method for globally solving nonconvex quadratic programming. *J Henan Norm Univ* 36(3):128–130
- Shen P, Chen Y, Ma Y (2009) A nonisolated optimal solution for special reverse convex programming problems. *J Comput Appl Math* 224:219–229
- Sherali HD, Smith EP (1997) A global optimization approach to a water distribution network design problem. *J Glob Optim* 11:107–132
- Thoi NV (2000) Duality bound method for the general quadratic programming problem with quadratic constraints. *J Optim Theory Appl* 107(2):331–354
- Vandenbussche D, Nemhauser GL (2005) A branch-and-cut algorithm for nonconvex quadratic programs with box constraints. *Math Program* 102:559–575
- Vavasis SA (1992) Approximate algorithms for indefinite quadratic programming. *Math Program* 57:279–311
- Voorhis TV (2002) A global optimization algorithm using lagrangian underestimates and the interval newton method. *J Glob Optim* 24:349–370

- Wang Y, Liang Z (2005) A deterministic global optimization algorithm for generalized geometric programming. *Appl Math Comput* 168:722–737
- Wang Y, Zhang KC, Gao LY (2004) Global optimization of generalized geometric programming. *Comput Math Appl* 48:1505–1516
- Yajima Y, Fujie T (1998) A polyhedral approach for nonconvex quadratic programming problems with box constraints. *J Glob Optim* 13:151–170
- Ye Y (1992) On the affine scaling algorithm for nonconvex quadratic programming. *Math Program* 56:285–300
- Zhang Y, Agarwal P, Bhatnagar V, Balochian S, Yan J (2013) Swarm intelligence and its applications. *Sci World J* 2013:1–3
- Zhang Y, Agarwal P, Bhatnagar V, Balochian S, Zhang X (2014a) Swarm intelligence and its applications 2014. *Sci World J* 2014:1–4
- Zhang Y, Balochian S, Agarwal P, Bhatnagar V, Housheya OJ (2014b) Artificial intelligence and its applications. *Math Probl Eng* 2014:1–10
- Zhang Y, Balochian S, Agarwal P, Bhatnagar V, Housheya OJ (2016) Artificial intelligence and its applications 2014. *Math Probl Eng* 2016:1–6

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com
